

TOWARDS SUCCESSFUL REINTRODUCTIONS: THE COMBINED IMPORTANCE
OF SPECIES TRAITS, SITE QUALITY, AND RESTORATION TECHNIQUE

Thomas N. Kaye

Institute for Applied Ecology, PO Box 2855, Corvallis, Oregon 97339-2855

Department of Botany and Plant Pathology, Oregon State University, Corvallis, Oregon

97331

ABSTRACT

Reintroduction of endangered species may be necessary to protect them from extinction, provide connectivity between populations, and reach recovery goals under the U.S. Endangered Species Act. But what factors affect reintroduction success? And which matter more, traits inherent to the species, qualities of the site, or reintroduction technique? Here I propose that all three interact. First, reintroduction success will be highest for endangered species that share traits with non-rare native species, invasive plants, and species that excel in restoration plantings as reviewed from the ecological literature. Ten traits are identified as common to at least two of these groups. Second, reintroductions will do best in habitats ecologically similar to existing wild populations and with few local threats, such as non-native plants and herbivores. And third, the methods used to establish plants, such as planting seeds vs. transplants or selecting appropriate microsites, will influence outcomes. For any reintroduction project, potential

pitfalls associated with a particular species, site or technique may be overcome by integrating information from all three areas. Conducting reintroductions as designed experiments that test clearly stated hypotheses will maximize the amount and quality of information gained from each project and support adaptive management.

Keywords: endangered species, extinction, plant trait, reintroduction, review

Kaye, T.N. In press. Towards successful reintroductions: The combined importance of species traits, site quality, and restoration technique. Proceedings of a conference on Strategies and Solutions for Plant Conservation. California Native Plant Society, Sacramento, CA.

INTRODUCTION

Endangered species reintroduction is necessary when too few populations exist in the wild to sustain long-term viability. It may be required to meet the objectives of Recovery Plans when their criteria call for additional populations in areas where a species has been extirpated. In addition, reintroduction may be implemented to mitigate for population losses caused by habitat development or changes in management priorities, but mitigation of this sort is much more controversial and fraught with ethical concerns (e.g., Allen 1994). In a review of 181 Recovery Plans for endangered species, one study (Hoekstra et al. 2002) found that 72% of plans call for some form of reintroduction. But reintroduction may be a costly process with no guarantee of success. Do some species lend themselves to reintroduction more easily than others?

Reintroduction is often regarded as a special form of habitat restoration that applies to rescuing or recovering endangered species (Maunder 1991; Falk et al. 1996; Armstrong & Seddon 2008). It faces many unique challenges due to the high value placed on the species targeted for improvement and the frequent knowledge gaps about individual species' needs (Guerrant & Kaye 2007). Reintroduction in general has been subject to such frequent failure that many regard it as unreliable (Fahselt 2007).

The reintroduction process has been outlined, evaluated, and updated over the last two decades by various workers around the world (e.g., Griffith et al. 1989, IUCN 1995, Vallee et al. 2004, Kaye 2008, Menges 2008). Falk et al. (1996) provide a

comprehensive overview of reintroduction. However, predicting which endangered species will perform best in a reintroduction has not been possible to date, partly because too few reintroductions have been published to draw broad generalizations.

Population reintroduction is a field still searching for a consistent vocabulary (Armstrong & Seddon 2008). Translocation is a term widely used for the same process, and can include the wholesale transplanting of individuals or populations from one wild site to another. Augmentation is the process of adding individuals to an existing population to increase its size and viability, and may be considered one form of reintroduction.

Introduction is sometimes used as a synonym for reintroduction or translocation, but also describes the process of non-native and invasive species movement into a new region. I use the term reintroduction here inclusively to include all forms of placing plant materials into occupied or unoccupied sites of an endangered species within its historic range or ecoregion, with the assumption that a species may have occurred in any piece of appropriate habitat at some point in the past even if there is no supporting historic documentation.

Over 740 plant taxa are listed by the US Fish and Wildlife Service as Threatened or Endangered (http://ecos.fws.gov/tess_public/SpeciesReport.do, accessed 5/24/09). With so many species needing support from conservationists, funding is unlikely to be available for a reintroduction program for all of them. Criteria for selecting species that are most likely to succeed in a reintroduction program are therefore urgently needed, and this will help maximize our efficiency. Also, species targeted for reintroduction that have

a low chance of success will need additional effort, and this effort will be assisted if we are able to anticipate and plan for problems.

In this paper, I suggest that three primary factors interact to control reintroduction success (defined here as establishment, growth and formation of a viable population, but see Pavlik [1996] for a comprehensive discussion of success in this context) and, when considered together, can improve reintroduction of rare species in general. These factors include plant traits inherent to the species, habitat conditions at the reintroduction site, and reintroduction technique. I consider each factor individually then conclude with a discussion of how they interact.

PLANT TRAITS

Plant traits have been used to predict species response to disturbance (McIntyre et al. 1995), competitive ability (Grace 1990, Goldberg 1996), and invasiveness (Baker 1974, Sakai et al. 2001), for example. Since plant traits may have predictive power, an evaluation of rare species based on specific traits may help identify species most suitable for reintroduction. In order to develop a list of likely traits for this purpose, I consulted reviews of plant traits in three topic areas based on the hypotheses that reintroduction success will be highest for rare species that share traits with:

1. non-rare native species (or have traits opposite of typical rare species),
2. invasive plants, and

3. species that excel in restoration plantings.

These three areas were selected because traits inherent to rarity may make species more vulnerable to rapid extirpation at a new location, species that share traits with invasive species may tend to do well and spread when placed at a new site, and rare plants most like those that succeed in restorations may do best at the establishment and colonization phase. I looked for plant traits common to at least two of these categories to develop a preliminary list of plant traits that identify species that are “most likely to succeed.”

A synthesis of reviews

Thorough reviews of the predictive power of plant traits in each of these topic areas are available (Table 1). For traits associated with rarity, I consulted three large reviews. Murray et al. (2002) searched for life history and ecological traits that consistently separate rare and common species. Their paper is really two papers in one; species in general from 54 comparative studies were examined from the global literature and 700 eucalypt species in Australia were evaluated from their own data. Farnsworth & Ogurcak (2008) examined 71 rare plant species in New England to determine if species with shared traits showed similar fates of range shift and population extirpation. Hedge & Ellstrand (1999) looked at the well-studied floras of California and the British Isles to identify life history differences between species known to be rare or common.

Two reviews by Kolar & Lodge (2001) and Hayes & Barry (2008) were used to identify traits of invasive species. In the first, the authors reviewed 16 publications (half on plants) on invasions to identify traits of successful invaders. The second paper added 33 more studies to those reviewed in the first to identify independently verified predictors of invasion or establishment success of non-native species across multiple biological groups, including plants.

Traits of plants that grow well in ecological restorations were identified by Pywell et al. (2003). They performed a meta-analysis of 25 experiments with a cumulative total of 58 species to identify traits that predicted performance in field restorations in agricultural lowland Britain. Forbs (45 species) and grasses (13 species) were analyzed separately.

A total of 38 plant traits were identified in this process that were significant in at least one review (Table 1). All of these traits may be considered hypotheses for characteristics of plants likely to do well when reintroduced. Traits were placed into categories of life history, plant size and growth, pollination, seed biology, habitat, history, and characteristics of the native range. Some reviews included additional traits that applied to other biological groups besides plants, and in some cases I combined multiple traits into a single trait for simplicity. Therefore, the traits in Table 1 reflect my interpretation, in some cases, of the traits identified by the various studies in an effort to reconcile differences in terminology while maintaining the authors' intent. A total of ten traits (listed in bold) were common to at least two categories. Overlap among groups provides

support for individual traits as predictors of reintroduction success because the overlap suggests some generality across contexts.

Traits of plants likely to succeed in a reintroduction

Taken together, the traits common to more than one category provide a starting point for a hypothetical list of characteristics of species that may be most adapted for reintroduction (Table 2). For example, plants with a life-history that includes a long flowering and/or fruiting period as well as vegetative reproduction or clonal growth may be good candidates for reintroduction because they are unlike most critically endangered species and share characteristics with invasives and species that do well in restorations. Plants that are competitive, tall and have large leaf area may perform well in reintroductions for similar reasons.

Plants with insect pollen vectors (as opposed to wind or self pollination) were both more likely to be rare and invasive. This apparent contradiction makes it difficult to identify a specific pollination mechanism as a reintroduction trait. Even so, a bet-hedging strategy for a reintroduction candidate would be a mixed-mating system based on generalist insect or wind pollination with self compatibility and autogamy as a failsafe mechanism.

High seed production is associated with invasiveness but not rarity, while small seed size is generally (but not always) associated with invasive species and restoration. There is a general ecological trade off between seed size, number, and seedling recruitment such

that seed size tends to decline as seed production increases, but species with larger seeds tend to establish and compete better than small seeded species (Jakobsson & Eriksson 2000). Species with intermediate seed weight may optimize seed production and seedling performance (Jakobsson & Eriksson 2000). Therefore, species that produce seeds of intermediate size in large numbers may stand the best chance of success during reintroduction. Another seed biology trait associated with reintroduction potential may be strong dispersal, since rare species tend to have poor dispersal mechanisms while invasives tend to be capable of long distance dispersal (Trakhtenbrot et al. 2005).

At least one analysis in each category found that geographic range size and/or variation in the types of habitats occupied by the species was an important factor in explaining species behavior. For example, rare species tend to occur over small geographic areas and occupy specialized habitats while invasives tend to have relatively large home ranges, and forbs that recruit well in restorations tend to be habitat generalists.

A history of invasion is a strong predictor of species invasiveness in new areas, as is taxonomic relationship to other invasives (Kolar & Lodge 2001, Hayes & Barry 2008). For most endangered species, no previous attempts at reintroduction have been performed, but this is likely to change with time as more reintroductions are attempted and the results published. I suggest that a history of reintroduction success or relationship (same genus or family) to species with a prior record of successful establishment will be a useful trait in identifying reintroduction projects with a high likelihood of success.

Climate change can affect native species in predictable ways. Species decline and loss near Concord, Massachusetts in Thoreau's woods was driven by climate change and the ability of species to adapt to temperature shifts (Willis et al. 2008). Species that persisted over the last 150 years as the climate warmed were those that tracked seasonal temperature and shifted their flowering times in response to year to year variation and long term trends. Plants that did not alter their phenology in response to the changing climate were more likely to become less abundant or die out. This response was shared among closely related taxa. Species that are able alter their flowering time as climate changes may have an advantage in reintroduction projects because climate already has changed in many areas and continues to change. Although a plastic phenology that shifts with climate was not identified as a significant trait in any of the studies reviewed in Table 1, its importance is a recent discovery and has obvious relevance to plant conservation. For that reason it is included among the traits of plants most likely to succeed in reintroductions (Table 2).

CHARACTERISTICS OF THE REINTRODUCTION SITE

Characteristics of the reintroduction site may be just as important as traits of the plant to be reintroduced. Reintroduction success may be highest in habitats ecologically similar to existing wild populations. For example, fitness of transplanted *Lotus scoparius* (Nutt.) Ottley was positively correlated with environmental similarity between the source population and the transplant site (Montalvo & Ellstrand 2000). Similarly, survival of

Castilleja levisecta Greenm., an endangered species of the Pacific Northwest, was highest at reintroduction sites most similar in vegetation to the seed source (Lawrence and Kaye 2009). In both cases, geographic distance between source and receptor site had little or no correlation with plant performance.

Non-native species may pose one of the most significant obstacles to reintroduction at any given location. Establishment and growth of *Abronia umbellata* ssp. *breviflora* (Standl.) Munz on Pacific Coast beaches was strongly negatively correlated with vegetative cover of *Ammophila arenaria* (L.) Link, a highly invasive grass (Kaye 2004). Weed competition adversely affected transplant survival and growth of two endangered Australian plants (Jusaitis 2005). Growth of planted *Castilleja levisecta* was negatively correlated with abundance of non-native plants across ten sites (Lawrence & Kaye 2009). Invasive species are clearly a threat to endangered plants, and without their control reintroductions are likely to have limited success in general.

Site productivity may also affect reintroduction success and interact with invasive species abundance. For example, high nutrient sites and areas that have received nutrient addition have lower diversity and increased dominance in wetlands (Bedford et al. 1999) and forests (Gilliam 2006). Invasive plants in California ecosystems are disproportionately more frequent in areas with high productivity, and the high frequency of rare and endangered species in these ecosystems makes them especially vulnerable to competition from invasives (Seabloom et al. 2006). Finally, nutrient addition into low productivity sites makes them more vulnerable to invasion by exotics (e.g., Lake &

Leishman 2004). Selecting sites with low productivity, or reducing productivity artificially, may improve the odds of reintroduction success, especially for endangered species that are typically non-dominant in their wild populations.

Removal of threats in general may be crucial for reintroduction success. Sites that have processes in place that are associated with the cause of a species' endangerment or inhibit major life history steps may be poor choices. Invasive weeds and changes increases in nutrient availability are obvious examples, but alterations of disturbance regimes in general, such as fire suppression (e.g., Kaye et al. 2001, Quintana-Ascencio et al. 2003, Menges & Quintana-Ascencio 2004), may threaten rare species. Herbivores also can suppress seedling recruitment (Maschinski et al. 2004) and transplant survival (Jusaitis 2005, Lawrence & Kaye 2008, 2009) of reintroduced species. Sites that are not managed for conservation or are subject to development may be threatened with alteration or destruction (i.e., habitat loss) in the long-term.

REINTRODUCTION TECHNIQUE

The methods used to establish plants at a site may affect project outcomes. Soil amendment and planting season can have strong effects on plant success. Adding fertilizer and planting in the fall resulted in only 3% and 18% survival of *Erigeron decumbens* Nutt. and *Horkelia congesta* Douglas ex Hook., respectively, after four years, while omitting fertilizer combined with spring planting yielded 48% and 84% survival (Kaye & Brandt 2005). Fertilizer increased plant size and fecundity in *Abronia*

umbellata transplanted to beaches, but had no effect on survival (Kaye 2004). In addition, fencing plants to protect them from herbivory can increase transplant survival (Wendelberger et al. 2007, Thorpe 2008).

Microsites selected for transplants can have a significant effect on transplant success (Jusaitis 2005). For example, gaps created by sod cutting increased survival of *Scorzonera humilis* L. at nutrient rich sites but decreased it at nutrient poor sites (Reckinger et al. 2009). Seeds of *Centaurea corymbosa* Pourr. planted in selected cliff microsites established plants with higher survival than seeds that fell at random in a wild population (Colas et al. 2008). Soil moisture in particular may be important for the establishment of some species (Maschinski et al. 2004). For example, topographic position in a restored wetland affected transplant survival of four rare species (Kaye and Brandt 2005).

Propagule type and size have been explored in a variety of studies, and in general, larger plants have better performance during reintroduction. Plants established as sown seeds tend to have lower survival than transplants (Guerrant 1996, Bowles et al. 1998, Jusaitis et al. 2004, Kaye 2004, Guerrant and Kaye 2007, Reckinger et al. 2009), but if seeds are plentiful low establishment may be acceptable if plug production costs are high (Kaye and Cramer 2003). Also, smaller transplants may perform poorly compared to larger transplants (Wendelberger et al. 2007). In some cases, however, plant size seems to matter little for growth or survival (Alley & Affolter 2004).

In general, attempting reintroductions with more than one technique may be an optimal approach. Implementing reintroductions as designed experiments can serve simultaneously as a way to compare techniques or test hypotheses and a bet-hedging strategy to maximize the likelihood that at least one method will have high success (Guerrant & Kaye 2007, Kaye 2008).

CONCLUSION

Improvement in the success rates of reintroductions is urgently needed to support conservation efforts for many rare species. Predicting success and anticipating challenges will make conservation more efficient. If species performance in reintroduction can be predicted then scarce funds and other resources can be allocated to those most likely to succeed. The plant traits identified here (Tables 1 and 2) as potential markers of species that may be pre-adapted for reintroduction should serve as a hypothesis to be tested when larger data sets on the results of many reintroduction projects are available. And until then, they may be used to assist with ranking species for reintroduction. But even if the traits hypothesized here accurately predict success, ranking species and funding recovery work based on these characteristics may not always be desirable. In fact, including traits that are opposite those of most rare plants, as done here, will leave out many endangered species in need of aggressive conservation.

I suggest that the three areas discussed in this paper interact to determine the results of reintroductions (Fig. 1). The importance of plant characteristics will depend on site

qualities and planting methods. For example, species with many positive traits for growth and survival may perform well even at poor sites and require the least amount of effort. But as the number of positive traits that a species possesses declines, site quality and technique will increase in importance. In cases where reintroduction will proceed despite a species' low predicted potential, managers may need to pick superior sites or prepare them better and/or put more effort into developing optimal planting practices.

Species conservation should emphasize protecting existing populations, but reintroduction is needed in some cases to stave off extinction and meet recovery goals.

ACKNOWLEDGMENTS

I would like to thank Bruce Pavlik for suggesting that I pursue this interesting topic, and for coordinating the symposium at which this paper was originally presented.

LITERATURE CITED

ALLEN, W. H. 1994. Reintroduction of endangered plants. *BioScience* **44**:65-68.

ALLEY, H. AND J. M. AFFOLTER. 2004. Experimental comparison of reintroduction methods for the endangered *Echinacea laevigata* (Boynton and Beadle) Blake. *Natural Areas Journal* **24**:345-350.

ARMSTRONG, D. P. AND P. J. SEDDON. 2008. Directions in reintroduction biology. *Trends in Ecology and Evolution* **23**:20-25.

BAKER, H. G. 1974. The evolution of weeds. *Annual Review of Ecology and Systematics* **5**:1-24.

BEDFORD, B. L., M. R. WALBRIDGE, A. ALDOUS. 1999. Patterns in nutrient availability and plant diversity of temperate North American wetlands. *Ecology* **80**:2151-2169.

BOWLES, M. L., J. L. MCBRIDE AND R. F. BETZ. 1998. Management and restoration ecology of the federally threatened Mead's milkweed, *Asclepias meadii* (Asclepiadaceae). *Annals of the Missouri Botanical Garden* **85**:110-125.

- COLAS, B., F. KIRCHNER, M. RIBA, I. OLIVIERI, A. MIGNOT, E. IMBERT, C. BELTRAME, D. CARBONELL AND H. FRÉVILLE. 2008. Restoration demography: a 10-year demographic comparison between introduced and natural populations of endemic *Centaurea corymbosa* (Asteraceae). *Journal of Applied Ecology* **45**:1468-1476.
- FAHSELT, D. 2007. Is transplantation an effective means of preserving vegetation? *Canadian Journal of Botany* **85**: 1007-1017.
- FALK, D. A., C. I. MILLAR AND M. OLWELL. 1996. Restoring Diversity: Strategies for Reintroduction of Endangered Plants. Island Press, Washington, D.C.
- FARNSWORTH, E. J. AND D. E. OGURCAK. 2008. Functional groups of rare plants differ in levels of imperilment. *American Journal of Botany* **95**:943-953.
- GILLIAM, F. S. 2006. Response of the herbaceous layer of forest systems to excess nitrogen deposition. *Journal of Ecology* **94**:1176–1191.
- GOLDBERG, D. E. 1996. Competitive ability: definitions, contingencies and correlated traits. *Philosophical Transactions of the Royal Society of London* **351**:1377-1385.
- GRACE, J. B. 1990. On the relationship between plant traits and competitive ability, pp. 51-65. In, J. B. Grace & D. Tilman [eds.], *Perspectives on Plant Competition*,. Academic Press, San Diego.

GRIFFITH, B., J. M. SCOTT, J. W. CARPENTER AND C. REED. 1989. Translocation as a species conservation tool: status and strategy. *Science* **245**: 477 – 480.

GUERRANT, E. O. 1996. Designing populations: demographic, genetic, and horticultural dimensions, pp.171–208. *In*, D. A. Falk, C. J. Millar, and M. Olwell [eds.], *Restoring Diversity*. Island Press, Washington, D.C.

GUERRANT, E. O. AND T. N. KAYE. 2007. Reintroduction of rare and endangered plants: Common factors, questions and approaches. *Australian Journal of Botany* **55**:362-270.

HAYES, K. R. AND S. C. BARRY. 2008. Are there any consistent predictors of invasion success? *Biological Invasions* **10**:483-506.

HEDGE, S. G. AND N. C. ELLSTRAND. 1999. Life history differences between rare and common flowering plant species of California and the British Isles. *International Journal of Plant Science* **160**:1083-1091.

HOEKSTRA, J. M., J. A. CLARK, W. F. FAGAN, AND P. D. BOERSMA. 2002. A comprehensive review of Endangered Species Act Recovery Plans. *Ecological Applications* **12**:630–640.

IUCN. 1995. IUCN/SSC Guidelines for re-introductions. Prepared by the SSC Re-introduction Specialist Group, Abu Dhabi, UAE.

JAKOBSSON, A. AND O. ERIKSSON. 2000. A comparative study of seed number, seed size, seedling size and recruitment in grassland plants. *Oikos* **88**:494-502.

JUSAITIS, M. 2005. Translocation trials confirm specific factors affecting the establishment of three endangered plant species. *Ecological Management and Restoration* **6**:61-67.

JUSAITIS, M., L. POLOMKAB AND B. SORENSEN. 2004. Habitat specificity, seed germination and experimental translocation of the endangered herb *Brachycome muelleri* (Asteraceae). *Biological Conservation* **116**:251-266.

KAYE, T. N. 2001. Common ground and controversy in native plant restoration: the SOMS debate, source distance, plant selections, and a restoration-oriented definition of native, pp. 5-12. *In*, R. Rose and D. Haase [eds.], Native Plant Propagation and Restoration Strategies. Nursery Technology Cooperative and Western Forestry and Conservation Association, Corvallis, Oregon.

KAYE, T. N. 2004. Reintroducing the endangered pink sand-verbena to Pacific Coast beaches: Direct seeding and out-planting, pp. 131-139. *In*, M.B. Brooks, S.K. Carothers, and T. LaBlanca [eds.], The Ecology and Management of Rare Plants of Northwestern

California: Proceedings from a 2002 Symposium of the North Coast Chapter of the California Native Plant Society. California Native Plant Society, Sacramento, California.

KAYE, T. N. 2008. Vital steps toward success of endangered plant reintroductions. *Native Plants Journal* **9**:313-322.

KAYE, T. N. AND A. BRANDT. 2005. Seeding and transplanting rare Willamette Valley prairie plants for population restoration. Unpublished technical report, Institute for Applied Ecology, Corvallis, Oregon.

KAYE, T. N. AND J. CRAMER. 2003. Direct seeding or transplanting: The cost of restoring populations of Kincaid's lupine (Oregon). *Ecological Restoration* **21**:224-225.

KOLAR, C. S. AND D. M. LODGE. 2001. Progress in invasion biology: predicting invaders. *Trends in Ecology and Evolution* **16**:199-204.

LAKE, J. C. AND M. R. LEISHMAN. 2004. Invasion success of exotic plants in natural ecosystems: the role of disturbance, plant attributes and freedom from herbivores. *Biological Conservation* **117**:215-226.

LAWRENCE, B. A. AND T. N. KAYE. 2008. Direct and indirect effects of host plants: Implications for reintroduction of an endangered hemiparasitic plant (*Castilleja levisecta*). *Madroño* **55**:151-158.

LAWRENCE, B. A. AND T. N. KAYE. 2009. Reintroduction of *Castilleja levisecta*: Effects of ecological similarity, source population genetics, and habitat quality. *Restoration Ecology*, in press.

MASCHINSKI, J., J. E. BAGGS, AND C. F. SACCHI. 2004. Seedling recruitment and survival of an endangered limestone endemic in its natural habitat and experimental reintroduction sites. *American Journal of Botany* **91**:691-698.

MAUNDER, M. 1992. Plant reintroduction: an overview. *Biodiversity and Conservation* **1**:51-61.

MCINTYRE, S., S. LAVOREL, AND R. M. TREMONT. 1995. Plant life-history attributes: their relationship to disturbance response in herbaceous vegetation. *Journal of Ecology* **83**:31-44.

MCLACHLAN, J.S., J. J. HELLMANN, AND M. W. SCHWARTZ. 2007. A framework for debate of assisted migration in an era of climate change. *Conservation Biology* **21**:297-302.

MENGES, E. S. 2008. Restoration demography and genetics of plants: When is a translocation successful? *Australian Journal of Botany* **56**, 187–196.

MENGES, E. S. AND P. F. QUINTANA-ASCENCIO. 2004. Population viability with fire in *Eryngium cuneifolium*: deciphering a decade of demographic data. *Ecological Monographs* **74**:79-99.

MONTALVO, A. M. AND N .C. ELLSTRAND. 2000. Transplantation of the subshrub *Lotus scoparius*: Testing the home-site advantage hypothesis. *Conservation Biology* **14** 1034-1045.

MURRAY, B., P. H. THRALL, A. M. GILL, AND A. B. NICOTRA. 2002. How plant life-history and ecological traits relate to species rarity and commonness at varying spatial scales. *Austral Ecology* **27**:291-310.

PAVLIK, B. 1996. Defining and measuring success, pp. 127-155. In, D. A. Falk, C. I. Millar, and M. Olwell [eds], *Restoring Diversity: Strategies for Reintroduction of Endangered Plants*. Island Press, Washington, D.C.

PYWELL, R. F., J. M. BULLOCK, D. B. ROY, L. WARMAN, K. J. WALKER, AND P. ROTHERY. 2003. Plant traits as predictors of performance in ecological restoration. *Journal of Applied Ecology* **40**:65-77.

QUINTANA-ASCENCIO, P. F., E. S. MENGES, AND C. W. WEEKLEY. 2003. A fire-explicit population viability analysis of *Hypericum cumulicola* in Florida rosemary scrub. *Conservation Biology* **17**:433-449.

RECKINGER, C., G. COLLING, AND D. MATTHIES. 2009. Restoring populations of the endangered plant *Scorzonera humilis*: Influence of site conditions, seed source, and plant stage. *Restoration Ecology*, in press.

SAKAI, A. K., F. W. ALLENDORF, J. S. HOLT, D. M. LODGE, J. MOLOFSKY, K. A. WITH, S. BAUGHMAN, R. J. CABIN, J. E. COHEN, N. C. ELLSTRAND, D. E.

MCCAULEY, P. O'NEIL, I. M. PARKER, J. N. THOMPSON, S. G. WELLER. 2001. The population biology of invasive species. *Annual Review of Ecology and Systematics* **32**:305-332

SEABLOOM, E. W. J. W. WILLIAMS, D. SLAYBACK, D. M. STOMS, J. H. VIERS AND A. P. DOBSON. 2006. Human impacts, plant invasion, and imperiled plant species in California. *Ecological Applications* **16**:1338-1350.

TRAKHTENBROT, A., R. NATHAN, G. PERRY AND D. M. RICHARDSON. 2005. The importance of long-distance dispersal in biodiversity conservation. *Diversity and Distributions* **11**:173–181.

THORPE, A. 2008. The good, the bad, and the ugly: Challenges in plant conservation in Oregon. *Native Plants Journal* **9**:351-357.

VALLEE, L., T. HOGBIN, L. MONKS, B. MAKINSON, M. MATTHES AND M. ROSSETTO. 2004. Guidelines for the translocation of threatened plants in Australia. 2nd edition. Australian Network for Plant Conservation, Canberra.

WENDELBERGER, K. S., M. Q. N. FELLOWS, J. MASCHINSKI. 2007. Rescue and restoration: Experimental translocation of *Amorpha herbacea* Walter var. *crenulata* (Rybd.) Isley into a novel urban habitat. *Restoration Ecology* **16**:542-552.

WILLIS, C. G., B. RUHFELA, R. B. PRIMACK, A. J. MILLER-RUSHING AND C. C. DAVIS. 2008. Phylogenetic patterns of species loss in Thoreau's woods are driven by climate change. *PNAS* **105**: 17029-17033.

.

Table 1. Plant traits of rare and endangered species, invasives, and top performers in restoration. Each column represents a review or synthesis of a large data set, or a subset of one. Only traits that were statistically significant in the original review are included. Traits in bold are common to at least two categories. The sign of the trait indicates the direction of effect. For example, rare species tend to have short flowering periods (-) but invasives tend to have long flowering periods (+). Traits with no sign for a review were not examined or were not significant in that review.

Trait	Rare & endangered			Invasive		Restoration	
	Citation	Murray et al. 2002	Farnsworth & Ogurcak 2008	Hedge & Ellstrand 1999	Kolar & Lodge 2001	Hayes & Barry 2008	Pywell et al. 2003
	General						
	Eucalyptus						
						forbs	grasses
Life history							
Length of flowering/seed production period		-	-		+	+	
Iteroporous				+			
Length of juvenile period					-	-	
Longevity						+	
Obligate symbioses			+				
Ruderality							+
Vegetative reproduction/clonal growth					+	+	+
Plant size and growth							

Taxonomic group			+	+	
Native range					
Range area/variation in habitats occupied	-		+	+	+
Geographic origin			+		
Match between habitat at source and destination				+	
At or near range limit		+			

Table 2. Traits of species likely to succeed in reintroduction.

1.	Long flowering/fruiting period
2.	Vegetative growth
3.	Competitive
4.	Tall
5.	Large leaf area
6.	Generalist pollination, mixed mating system
7.	High seed production, intermediate seed size
8.	Long distance dispersal mechanism
9.	Large geographic range and/or multiple habitats
10.	History of reintroduction success or taxonomic relationship to species with a history of success.
11.	Plastic phenology that shifts with climate




Figure 1. Reintroduction potential of endangered species is affected by species traits, site characteristics, and reintroduction technique. Species without traits that improve their likelihood of reintroduction success may need better sites or more effort toward cultivation or planting methods.