

LOMATIUM COOKII POPULATION MONITORING IN THE ILLINOIS VALLEY, JOSEPHINE COUNTY, OREGON



2019

Report to the Bureau of Land Management,
Medford District

Report prepared by Ian A. Pfingsten, Thomas N.
Kaye and Denise E.L. Giles
Institute for Applied Ecology



PREFACE

This report is the result of a cooperative project between the Institute for Applied Ecology (IAE) and the Bureau of Land Management. IAE provides services to public and private agencies and individuals through development and communication of information on ecosystems, species, and effective management strategies. Restoration of habitats, with a concentration on rare and invasive species, is a primary focus. IAE conducts its work through partnerships with a diverse group of agencies, organizations and the private sector. IAE aims to link its community with native habitats through education and outreach.



Questions regarding this report or IAE should be directed to:

Thomas Kaye (Executive Director)

Institute for Applied Ecology

563 SW Jefferson Avenue

Corvallis, Oregon 97333

phone: 541-753-3099

email: tom@appliedeco.org

ACKNOWLEDGMENTS

We thank the Medford BLM, particularly Bryan Wender and Rachel Showalter, for their support and assistance with this project. The following IAE staff and interns contributed to the 2019 field season: Michel Wiman, Jenn Thornhill, Miranda Geller, Eva Brod, and Cia Crowe.

Cover photograph: *Lomatium cookii* (Cook's Desert Parsley), and Illinois Valley habitat, Josephine County, Oregon.

Suggested Citation

Pfingsten, I.A., T.N. Kaye, and D.E.L. Giles. 2019. *Lomatium cookii* population monitoring in the Illinois Valley, Josephine County, Oregon. Institute for Applied Ecology, Corvallis, Oregon and USDI Bureau of Land Management, Medford District. ix+44 pp.

TABLE OF CONTENTS

PREFACE.....	II
ACKNOWLEDGMENTS.....	III
LIST OF FIGURES	VI
LIST OF TABLES	VIII
EXECUTIVE SUMMARY.....	IX
INTRODUCTION	1
Project Overview	1
Review of past monitoring	1
Description.....	2
Geographic range	2
Habitat.....	2
Reproductive biology.....	2
Concerns	2
Objectives	3
METHODS.....	5
Review of populations	5
French Flat Middle and South subpopulations	5
Middle subpopulation	7
South subpopulation	9
Density plots.....	10
Demographic plots and analysis.....	15
Rough and Ready Creek.....	18
Indian Hill	22
RESULTS.....	24
French Flat.....	24
Population size and density	24
Population structure	24
Demography and Population Viability Analysis.....	30
Rough and Ready Creek.....	34
Indian Hill	37
Herbivory on <i>Lomatium cookii</i>	39
DISCUSSION	41
Current population status	41
French Flat	41
Rough and Ready Creek	41
Indian Hill.....	41
Management Recommendations	42
French Flat	42
Rough and Ready Creek	43

LOMATIUM COOKII POPULATION MONITORING IN THE ILLINOIS VALLEY, JOSEPHINE COUNTY, OREGON

Indian Hill..... 43

LITERATURE CITED44

APPENDIX A. GEAR LIST.....46

LIST OF FIGURES

Figure 1. Line drawing of <i>Lomatium cookii</i> from Kagan (1986).	1
Figure 2. Locations of the <i>Lomatium cookii</i> study sites included in this report. Note: other <i>L. cookii</i> subpopulations occur in this general area but are not shown.....	4
Figure 3. Map of three French Flat subpopulations of <i>Lomatium cookii</i> . The Middle and South subpopulations include long-term monitoring subplots and transects, while the smaller North subpopulation is sampled with radially-arranged transects only.	6
Figure 4. Layout of new density and demography plots established in 2012 at French Flat South and in 2013 at Middle. Density plots run perpendicular to the baseline to the east or west, and are monitored in 10 cm strips on the south side of the transect line. Demography plots are marked in the northeast and southwest corners (NE is tagged).....	12
Figure 5. Location of French Flat Middle demographic and density plots established in 2013. Baseline transect is 100 m with a bearing of 034° (northeast). Last plants located on density plots are indicated by red crosses. Demography plots are indicated by blue squares and are not located along the baseline transect as shown here. See Table 1 for exact locations of all density and demography plots. Note that this figure does not reflect the transects on the east side of the baseline were extended in 2018.	13
Figure 6. Location of French Flat South demographic and density plots established in 2012. Baseline transect is 157 m with a bearing due south. Last plants located on density plots are indicated by red crosses. Demography plots are indicated by blue squares. See Table 2 for exact locations of all density and demography plots.....	14
Figure 7. Layout of new density and demography plots established in 2012 at French Flat South. Baseline transect is 157 m long. Density plots are run perpendicular to the baseline to the east or west. Density plots are monitored in 10 cm strips on the south side of the transect line. Demography plots are marked in the northeast and southwest corners (NE is tagged.) A table with locations of all density and demography plot locations can be found in Table 2.	19
Figure 8. Sketch map of the <i>L. cookii</i> patches located at Rough and Ready Botanical Wayside. Cross-hatching indicates patches of plants. The long-term monitoring transects are also indicated in the drawing. Note that the patches are distinguished by the letters A-H.....	20
Figure 9. Orientation of the sampling plots around transects 1, 2 and 3 at the Rough and Ready Botanical Wayside population.	21
Figure 10. <i>L. cookii</i> (left), and <i>L. utriculatum</i> (right). These species can co-occur in sampling plots.	22
Figure 11. Plot layout of the <i>Lomatium cookii</i> population at Indian Hill. Sample plots are 50 meters x 0.5 meters, marked at either end by a rebar post (with a copper tag and flagging). The plots are monitored on the east side of the tape.	23
Figure 12. The total number of <i>Lomatium cookii</i> at Middle and South French Flat, 1993-2019. Error bars represent a 95% confidence interval.....	25
Figure 13. The total number of reproductive <i>Lomatium cookii</i> at Middle and South French Flat, 1993-2019. Error bars represent a 95% confidence interval.....	25
Figure 14. The number of reproductive <i>Lomatium cookii</i> relative to the total subpopulation at Middle and South French Flat, 1993-2019.....	26
Figure 15. <i>Lomatium cookii</i> population structure in 2019 (top) and averaged over the course of the study (bottom) for the Middle French Flat subpopulation (1994-2019), South French Flat subpopulation (1994-2019), Rough and Ready Creek (1995-2019), and Indian Hill (1997-2019). Bars indicate the mean percentage of plants in each stage. Error bars in the lower graph represent +1 SE. See text for explanation of life-history stages.	29

Figure 16. Population trends and the total number of plants (reproductive and vegetative) at Rough and Ready Creek, 1994-2019. There were only 26 reproductive plants recorded in 1994.....	36
Figure 17. Population fluctuations in <i>L. cookii</i> patches A-H at Rough and Ready Creek, 1994-2019. Patches B-F were combined in 2015 and patches E-F were combined in 2018.....	36
Figure 18. Number of vegetative and reproductive plants at Indian Hill, 1997-2019.....	37
Figure 19. Frequency of herbivory on <i>L. cookii</i> populations at four populations, 1994-2019. Error bars represent 95% confidence intervals. All plants were measured at Rough and Ready population.....	40
Figure 20. ORV damage in the <i>Lomatium cookii</i> population at French Flat.	42

LIST OF TABLES

Table 1. Plot information for density and demography plots established in 2013 and 2019 at French Flat Middle.	8
Table 2. Plot information for density and demography plots established in 2012 at French Flat South. .	10
Table 3. Summary of <i>Lomatium cookii</i> population data for the South and Middle French Flat subpopulations, 1993-2019. Population size and density estimates include all life-history stages from seedlings to large reproductive plants.....	27
Table 4. Average transition matrices used in the stochastic analyses for <i>L. cookii</i> at French Flat. The top row of each matrix represents the number of seedlings produced per reproductive plant (fecundity). Lower rows represent the probability of an individual in a given stage changing to another stage the following year. Stage-specific mortality is provided in the bottom row. The elements in the matrices are averaged from the transition data collected in 1994-2011 and 2012-19 for the South subpopulation and 1994-99 and 2008-19 for the Middle subpopulation.	31
Table 5. Population growth and mortality rates at the French Flat Middle and South sub-populations from 1994-2019. [Note: Demographic data was not collected at the Middle subpopulation from 2000 through 2007, and plants mapped prior to 2012 were not relocated at the South subpopulation.]	32
Table 6. Elasticity matrices of the south and middle <i>Lomatium cookii</i> subpopulations derived from the average transition matrices. Elasticities represent the relative sensitivity of the population growth rate (λ) to small changes in the transition probabilities. The sum of all elasticities is 1.0, and the sum of each column is the total proportional sensitivity of λ to changes in transition probabilities in that stage. The top two individual elasticities are shown in bold.....	33
Table 7. Number of <i>L. cookii</i> individuals in each patch at Rough and Ready Creek. The total number of plants in all patches combined represents a census of the population. Totals include plants in all stages from seedlings to large reproductive plants. In 2015, patches C, D, E, and F were lumped into patch B, and in 2018, patch E was lumped into F.	35
Table 8. Summary of population density (mean $\pm$ 1 S.E.) and size (mean $\pm$ 95% confidence limit) for the Indian Hill population, 1997-2019.....	38

EXECUTIVE SUMMARY

Lomatium cookii is endemic to southwestern Oregon, and is listed as endangered by the State of Oregon and by the U.S. Fish and Wildlife Service (USFWS). Populations discussed in this report are located in the Illinois Valley in Josephine County. Long-term monitoring has occurred at French Flat since 1990, at Rough and Ready since 1994 and at Indian Hill since 1997. Data from these efforts are used to track population dynamics at all sites and create a population viability analysis for French Flat.

French Flat South

- Total population size was estimated at 50,811 individuals in 2019, a 20% decrease from 2018. Average plant density was estimated at 6.4 plants m⁻². The proportion of reproductive plants in the overall population was 33%, a 25% increase from 2018.
- The Population Viability Analysis (PVA) indicated a 99.9% probability this population will decrease 50% in the next 20 years. A catastrophic loss (99% population decrease) over the same period is less likely to occur (1.9%).

French Flat Middle

- The total population numbered 41,600 in 2019, an 8% increase from 2018. Density was estimated at 8.9 plants m⁻². Reproductive plants composed just over 24% of the overall population, a 29% decrease from 2018.
- The PVA indicated a 99.7% probability this population will decrease 50% in the next 20 years. A catastrophic loss (99% population decrease) over the same period has an approximately 13% probability to occur.

Rough and Ready

- This population increased 50% from 2018 to 2,444 plants in 2019. Reproductive adults comprised 39% of the population, a 15% decrease from 2018. Barricading with boulders has successfully blocked vehicles from dumping trash in the small meadow and accessing side roads.

Indian Hill

- This population increased 2% to 8,225 in 2019. Reproductive adults comprised 44% of the population, a 27% increase from 2018.

Overall, populations of *Lomatium cookii* in Josephine County have shown increased recruitment of seedlings and population sizes have remained relatively stable. This is likely due in part to active management of these populations by the Medford BLM, including preventing use by off road vehicles. However, population viability analyses indicate there is still risk of serious population declines. Proposed mining activities may have further detrimental impacts on these populations at French Flat. We recommend continued monitoring of these populations to assess population trends, trespass by ORVs, and provide data to assess future impacts to populations from climate change, mining, or other threats.

LOMATIUM COOKII POPULATION MONITORING IN THE ILLINOIS VALLEY, JOSEPHINE COUNTY, OREGON

REPORT TO THE BUREAU OF LAND MANAGEMENT, MEDFORD DISTRICT

INTRODUCTION

Project Overview

Lomatium cookii (cover photo,





Herbivory by voles is relatively frequent on *L. cookii*, but the frequency of damage to the plants has varied substantially among sites and years. Typically, damage to the plants ranges from consumption of leaf tips and inflorescences, to removal of nearly all aboveground plant material. Consumption of the flower clusters results in little damage to the individual plant, but eliminates seed production and, therefore, recruitment of seedlings in the following year. The long-term impact of herbivory on *L. cookii* plants is unknown, but appears to play an important role in population growth and seedling dynamics. Environmental parameters (e.g., rainfall, average maximum daily temperature, and average minimum daily temperature) may influence vole populations similarly across the region and in turn affect the number of *L. cookii* that are grazed.

Herbivory on *L. cookii* has fluctuated between years, and the patterns are not synchronous among the sites (Figure 19). Average herbivory at French Flat Middle and South increased since initial monitoring. In 2019, French Flat South and Rough and Ready populations had decreased herbivory from 2018, while French Flat Middle and Indian Hill populations had increased herbivory in the same year. In our observations since 1993, the highest recorded average herbivory was 57% (95% C.I.: 46% - 68%) detected at Indian Hill in 2019, while the lowest was 1% (95% C.I.: 0% - 2%) at French Flat Middle in 1994.



Figure 1. Line drawing of *Lomatium cookii* from Kagan (1986).

LOMATIUM COOKII POPULATION MONITORING IN THE ILLINOIS VALLEY, JOSEPHINE COUNTY, OREGON



DISCUSSION

Current population status

In 2019, the number of *L. cookii* increased at all sites except for French Flat South.

French Flat

Current population estimates at this site are approximately 41,600 plants at the Middle subpopulation and 50,811 at the South subpopulation. The population viability analysis suggests that the South and Middle subpopulations may be at risk of long term decline if current conditions persist. ORV use remains frequent and damaging (Figure 20). IAE is analyzing the effects of climate on *L. cookii* through Climate Driven Population Models and the results will be included in a separate report.

The Middle subpopulation hosted an estimated 195,057 individuals in 1998 when the population was at its highest, up from 74,694 when monitoring began in 1993 and over seven times the current estimate. Although there has been some variability, the population has generally been declining since 2000. The South subpopulation also peaked in 1998, with a total of 267,075 individuals. Since then, it has generally declined as well.

Results from population viability analyses based on information from 1994-99 and 2008-19 for Middle and 1994-2011 and 2012-19 for South indicated that these subpopulations differed in their long-term growth rates by about 2%, although both are likely to decline. Stochastic simulations based on observed environmental variability indicated that the risk of extinction or catastrophic decline was high at the Middle and South subpopulation if past environmental and biological conditions persist.

Rough and Ready Creek

We counted 2,444 plants at Rough and Ready Creek in 2019, which was a 50% increase from 2018. The Rough and Ready Creek population occurs in grassy patches and roadways on gentle to moderate slopes among shrubs and trees (*Pinus jeffreyi*). This patchy, less extensive habitat pattern is probably responsible for the large difference in population size between this site and French Flat. While road use can reduce the number of plants in this population, we observed recruitment along the roadbeds in 1995-2006, likely due to the reduced competition in these areas. A large trash pile found on the western edge of Patch A in 2003 and fresh ORV tracks in multiple years are indicative of active road use and off-road habitat impact. Boulders placed to block vehicle traffic on some roads and grassland patches has resulted in effective protection of this habitat, and grasses are now colonizing the protected road beds.

Indian Hill

The Indian Hill population, which is intermediate in size between the Rough and Ready and French Flat populations, declined in total numbers from 1997 (when monitoring began) to 1998, then remained relatively stable from 1998 to 2004. Overall, the population has increased slightly and remained stable from 2005-19, although 2016 had the second lowest population size on record. Estimates of total plant density and reproductive plant density followed similar changes. Many of the V3 and R1 plants are likely a result of the prolific seedling recruitment in 2005 and 2011. The proportion of seedlings in the population has been increasing since a low point in 2014. The population size has been relatively stable in recent years, and the site does not appear to receive ORV use. The error around our estimates of plot density and population size at Indian Hill have increased since monitoring began. This is due to high variability among transects and increased shrub density from *Ceanothus cuneatus*.



Figure 20. ORV damage in the *Lomatium cookii* population at French Flat.), Cook's desert-parsley, is listed as endangered by the State of Oregon and by the U.S. Fish and Wildlife Service (USFWS). It is closely related to *L. bradshawii* (Rose) Math. & Const., an endangered species found in the Willamette Valley of western Oregon. Several significant populations of *L. cookii* occur on land managed by the Bureau of Land Management (BLM) Medford District. This progress report documents monitoring methods and results on the BLM Medford District through 2019 and is the product of a cooperative effort between the BLM and the Institute for Applied Ecology.

Review of past monitoring

The largest population of *L. cookii* on federal land was recorded in 1992 at French Flat on the BLM Medford District. Areas around this population were placer-mined for many years. During the 1993 field season, staff from the BLM Medford District established long-term monitoring plots and transects in the three largest subpopulations of *L. cookii* in this area (Tong 1993). Additional plots and documentation of the monitoring protocol were added in 1994 (Kaye and Kirkland 1995). A long-term monitoring transect located southeast of these subpopulations was established by BLM Medford District botanists in 1990 (Knight 1992) and was monitored through 1994 (Kaye and Kirkland 1995), but has not been relocated in recent years and is not discussed in this report. Annual monitoring plots were

established on BLM land near Rough and Ready Creek in 1994 and at Indian Hill in 1997. Monitoring between 1994 and 1999 was conducted by the Oregon Department of Agriculture Native Plant Conservation Program. Since 2000, populations have been monitored by the Institute for Applied Ecology.





Description

***L. cookii* is a member of the Apiaceae (parsley family). The plants are usually less than 3 dm tall and inconspicuous except when in flower. Ternately divided leaves feature many narrow leaflets and creamy yellow flowers are produced in compound umbels on leafless stems.**

Herbivory by voles is relatively frequent on *L. cookii*, but the frequency of damage to the plants has varied substantially among sites and years. Typically, damage to the plants ranges from consumption of leaf tips and inflorescences, to removal of nearly all aboveground plant material. Consumption of the flower clusters results in little damage to the individual plant, but eliminates seed production and, therefore, recruitment of seedlings in the following year. The long-term impact of herbivory on *L. cookii* plants is unknown, but appears to play an important role in population growth and seedling dynamics. Environmental parameters (e.g., rainfall, average maximum daily temperature, and average minimum daily temperature) may influence vole populations similarly across the region and in turn affect the number of *L. cookii* that are grazed.

Herbivory on *L. cookii* has fluctuated between years, and the patterns are not synchronous among the sites (Figure 19). Average herbivory at French Flat Middle and South increased since initial monitoring. In 2019, French Flat South and Rough and Ready populations had decreased herbivory from 2018, while French Flat Middle and Indian Hill populations had increased herbivory in the same year. In our observations since 1993, the highest recorded average herbivory was 57% (95% C.I.: 46% - 68%) detected at Indian Hill in 2019, while the lowest was 1% (95% C.I.: 0% - 2%) at French Flat Middle in 1994.

LOMATIUM COOKII POPULATION MONITORING IN THE ILLINOIS VALLEY, JOSEPHINE COUNTY, OREGON



DISCUSSION

Current population status

In 2019, the number of *L. cookii* increased at all sites except for French Flat South.

French Flat

Current population estimates at this site are approximately 41,600 plants at the Middle subpopulation and 50,811 at the South subpopulation. The population viability analysis suggests that the South and Middle subpopulations may be at risk of long term decline if current conditions persist. ORV use remains frequent and damaging (Figure 20). IAE is analyzing the effects of climate on *L. cookii* through Climate Driven Population Models and the results will be included in a separate report.

The Middle subpopulation hosted an estimated 195,057 individuals in 1998 when the population was at its highest, up from 74,694 when monitoring began in 1993 and over seven times the current estimate. Although there has been some variability, the population has generally been declining since 2000. The South subpopulation also peaked in 1998, with a total of 267,075 individuals. Since then, it has generally declined as well.

Results from population viability analyses based on information from 1994-99 and 2008-19 for Middle and 1994-2011 and 2012-19 for South indicated that these subpopulations differed in their long-term growth rates by about 2%, although both are likely to decline. Stochastic simulations based on observed environmental variability indicated that the risk of extinction or catastrophic decline was high at the Middle and South subpopulation if past environmental and biological conditions persist.

Rough and Ready Creek

We counted 2,444 plants at Rough and Ready Creek in 2019, which was a 50% increase from 2018. The Rough and Ready Creek population occurs in grassy patches and roadways on gentle to moderate slopes among shrubs and trees (*Pinus jeffreyi*). This patchy, less extensive habitat pattern is probably responsible for the large difference in population size between this site and French Flat. While road use can reduce the number of plants in this population, we observed recruitment along the roadbeds in 1995-2006, likely due to the reduced competition in these areas. A large trash pile found on the western edge of Patch A in 2003 and fresh ORV tracks in multiple years are indicative of active road use and off-road habitat impact. Boulders placed to block vehicle traffic on some roads and grassland patches has resulted in effective protection of this habitat, and grasses are now colonizing the protected road beds.

Indian Hill

The Indian Hill population, which is intermediate in size between the Rough and Ready and French Flat populations, declined in total numbers from 1997 (when monitoring began) to 1998, then remained relatively stable from 1998 to 2004. Overall, the population has increased slightly and remained stable from 2005-19, although 2016 had the second lowest population size on record. Estimates of total plant density and reproductive plant density followed similar changes. Many of the V3 and R1 plants are likely a result of the prolific seedling recruitment in 2005 and 2011. The proportion of seedlings in the population has been increasing since a low point in 2014. The population size has been relatively stable in recent years, and the site does not appear to receive ORV use. The error around our estimates of plot density and population size at Indian Hill have increased since monitoring began. This is due to high variability among transects and increased shrub density from *Ceanothus cuneatus*.



Figure 20. ORV damage in the *Lomatium cookii* population at French Flat.). Fruits are flattened and oblong. The species was originally described by Kagan in 1986 from specimens collected in the Medford area.

Geographic range

L. cookii is endemic to southwestern Oregon. Two population centers are known, the Illinois Valley in Josephine County and the Agate Desert north of the Medford Plains in Jackson County (Kagan 1994). This report focuses on population monitoring on BLM land in the Illinois Valley at French Flat, Indian Hill, and Rough and Ready Creek (Figure 2). The Nature Conservancy currently monitors populations on their lands in Jackson County.

Habitat

The populations of *L. cookii* studied in the Illinois Valley are found in moist, grassy meadows dominated by *Danthonia californica* (Kaye and Blakeley-Smith 2002). Other associated species include *Deschampsia cespitosa*, *Festuca roemerii* ssp. *klamathense*, *Stipa lemmonii*, *Camassia quamash*, *Ranunculus occidentalis*, *Hesperochiron occidentalis*, *Downingia yina*, *Horkelia daucifolia*, *Isoetes nuttallii*, *Calochortus uniflorus*, and

Viola hallii. One patch of *L. cookii* individuals at the Rough and Ready Botanical Wayside was observed growing up through a dense ground-cover of *Toxicodendron diversilobum*. Trees and shrubs, such as *Pinus ponderosa*, *P. jeffreyi*, *Arctostaphylos* spp., and *Ceanothus cuneatus* border these grassy meadows.

Reproductive biology

Flowering stems begin to emerge from a rosette of leaves in late February and flowers usually bloom around mid-March and continue into May. As with some other *Lomatium* species, the earliest umbels are predominately staminate, while later umbels have both staminate and hermaphroditic flowers. Plants that produce only one umbel produce very few, if any, fruits (Kaye and Kirkland 1994). Several pollinators have been observed visiting *L. cookii* including a small bee in the Andrenae family (Brock 1987) and a small black moth (Kagan 1986). During 1994 and 1995, we observed large numbers of bumblebees (*Bombus* spp.) regularly visiting the flowers at French Flat in the Illinois Valley.

Concerns

Mining activities continue to threaten *L. cookii*. Placer gold mining has restricted the population at French Flat and permanently altered much of the natural hydrologic patterns through the meadows. Hydrology at the Rough and Ready Creek population is affected by a nearby irrigation ditch and by roads that pass through and divide the population. Some of the French Flat subpopulations monitored and discussed in this report are located on BLM managed lands adjacent to the Hillside Placer No. 1 and No. 3 Mines owned and operated by a local resident. A proposed mining plan filed in 1993 would involve destruction of a significant portion of this subpopulation. Recently, mining plans have been filed with BLM that will alter habitat immediately adjacent to *L. cookii* at French Flat ACEC.

L. cookii habitat in the Illinois Valley is threatened by rural development and abuse by recreational users in the area. Both the French Flat and Rough and Ready Creek sites continue to be severely damaged by ORV use, where we observed fresh vehicle tracks in 2002-2007. At both sites, damage was severe in the *L. cookii* population, even disturbing population monitoring plot markers. A trash pile, with old appliances and deep tire ruts, was found at the Rough and Ready Creek population in 2003. Tire ruts were found again at site in 2007.

Objectives

The purpose of this cooperative project is to assist in recovery efforts for *L. cookii* through the following population monitoring tasks:

- Resample existing monitoring plots on BLM land near French Flat to monitor changes in population size and/or density.
- Resample existing monitoring plots at the Rough and Ready Creek Botanical Wayside to monitor changes in population size.
- Resample permanent monitoring plots at the Indian Hill population.
- Summarize monitoring data from 1993 through 2019.
- Summarize information from long-term demographic plots within the French Flat subpopulation to provide information on the mortality and survival of plants in six life-history stages and develop a population viability model based on information from 1994-2019.

Image removed from web version

Figure 2. Locations of the *Lomatium cookii* study sites included in this report. Note: other *L. cookii* subpopulations occur in this general area but are not shown.

METHODS

Review of populations

Study populations included in this report are French Flat, Rough and Ready Creek, and Indian Hill. Monitoring has been conducted at four subpopulations, North, Middle, South, and Piney Woods (Figure 2), in or near French Flat since 1990. Currently, only the middle and south subpopulations (Figure 3) are monitored annually. Monitoring at Piney Woods was initiated in 1990 (Knight 1992) and conducted in 1994, but is not discussed here. Transects and monitoring at French Flat North was initiated by BLM staff in 1993 and was repeated in 1994 (Tong 1993), but is not discussed here. Transect and monitoring plots were also initiated by BLM staff at French Flat South and Middle in 1993 (Tong 1993). Sample plot numbers and sizes were modified in 1994 and the updated protocols are described below. These subpopulations continue to be monitored annually. Monitoring plots at Rough and Ready Creek were initiated in 1994. At Indian Hill, plots were established and sampled in 1997. The monitoring and sampling methods for each study site differ significantly, and are described separately. Units of measure (meters or feet) differ because different people initiated the monitoring plots at specific sites.

French Flat Middle and South subpopulations

Long-term monitoring plots were established at the Middle and South subpopulations in 1993. Density monitoring plots (1 foot x 10 feet and 1 foot x 20 feet) were located randomly within x, y coordinate grids positioned to cover the entire area of each subpopulation. The origins of the grid baselines were permanently marked with rebar. During the 1994 season, we recorded x, y coordinates for each of the plots at these subpopulations, measured locations and bearings of reference transects, added additional sampling plots, and adjusted the plot sizes [see discussions for each subpopulation below, and Kaye and Kirkland (1995)].

Two corners of each plot were permanently marked with rebar and metal pins were temporarily used for the other two corners of the rectangular plots during sampling. At some point between the 2011 and 2012 field season, all plot markers were removed from the French Flat South subpopulation. Only 3 of the 44 rebar remained in place and the plots could not be re-established. New plots were established in 2012 at South and in 2013 at Middle (see below for details).

Image removed from web version

Figure 3. Map of three French Flat subpopulations of *Lomatium cookii*. The Middle and South subpopulations include long-term monitoring subplots and transects, while the smaller North subpopulation is sampled with radially-arranged transects only.

Middle subpopulation

1993-2012

To locate the baseline grid in the Middle subpopulation, a tape representing the y-axis was stretched at a bearing of 035° east from the origin rebar (marked with a numbered copper tag and flagged) to a rebar post at 165 feet marked with a BLM "vegetation study" sign. Forty plots were established at random locations within the grid system. Demographic plots were established at sampling plots numbered 1, 3, 6, 7, 8, 10, 13, 16, 17, 18, 23, 25, 28, 29, 30, 31, 33, 37, 38, and 39.

2013-2019

New density plots were established in 2013 to reflect the methods at French Flat South in 2012 (see below). The boundary of the population was mapped, and a 100 m baseline transect was laid at a bearing of 034° (northeast) to bisect the population. Thirty density plots were placed at random intervals perpendicular (either east or west) to the baseline transect (Table 1, Figure 4, and Figure 5). A tape was run for 40 m and a 10 cm strip on the south side of the tape was monitored for *L. cookii*. Plants in the density plots were assigned to the life-history categories detailed below.

Previous demography plots were located. Demography plots without plants for multiple years were abandoned and four new plots were established. Only two demography plots were not relocated in 2013. All demographic plots monitored since 2014 were relocated and monitored.

After burning and meadow clearing at French Flat in 2017 the population has expanded such that some of our original transects at French Flat Middle did not capture the whole population. Transects 155, 161, 163, 167, 30, 171, 35, 173, 159, 177, and 179 were extended to 60 m on the east side of the baseline transect, while transect 195 was extended to 33.3 m on the east side as well (Table 1).

LOMATIUM COOKII POPULATION MONITORING IN THE ILLINOIS VALLEY, JOSEPHINE COUNTY, OREGON

Table 1. Plot information for density and demography plots established in 2013 and 2019 at French Flat Middle.

2013 New Plot #	Side of Tape	Location on Baseline (m)	End Rebar at (m)	End Rebar Tag	2013 Last Plant Found at (m)	Demog. Tag	Demog. Plot Location (NE Corner)	Demog. Plot Location (End)
165	W	2.8	30	166	25.95	33	11.43	10.90
154	W	6.5	37	153	34	1	11.2	10.75
155	E	6.5	60	156	26.5	3	18.55	18.05
161	E	9.1	60	162	35.2	8	23.25	23.75
163	E	9.7	60	164	38.05	7,10	7: 15.5; 10: 33.25	7: 16; 8: 33.75
28	W	15	37	29	35.7	876	18.55	19.05
167	E	17.5	60	168	38.05	-	-	-
30	E	19	60	31	35.1	-	-	-
33	W	22	40	34	28.1	877	23.45	23.95
169	W	25	40	170	38.7	37	22.24	21.71
171	E	27.1	60	172	33	38	2.4	2.9
199	W	31	30	200	29.3	-	-	-
35	E	35	60	36	38.6	-	-	-
173	E	36.6	60	174	36.6	874,875	874: 20.5; 875: 11.2	874: 30; 875: 11.7
175	W	40.2	35	176	31.85	167	23.45	23.95
158	W	43.1	30	157	20.6	18	13.2	12.7
159	E	43.5	60	160	28.25	28	5.2	5.7
177	E	46.6	60	178	33.9	29	5.1	5.6
179	E	55.1	60	180	13.7	31	4.75	5.25
181	W	56	20.4	182	19	-	-	-
183	E	60	11.3	184	11.05	-	-	-
185	E	62	10.8	186	9.7	-	-	-
187	W	67	9.9	188	6.7	-	-	-
189	E	72	15	190	14.9	168	6.5	7
37	W	74	16	38	6.9	-	-	-
191	W	82	15.5	192	11.4	169	2.5	2
193	W	86	10.5	194	3.5	-	-	-
195	E	89	33.3	196	19.5	170	15.6	15.1
197	W	95	15	198	10.5	-	-	-
39	W	98	15	40	8.7	-	-	-

South subpopulation

1993-2011

Foot/meter tapes attached end-to-end representing the y-axis were stretched at a bearing of 190° east from the origin rebar to a rebar post approximately 400 feet away. A numbered copper tag was attached to the northeast corner rebar post of each plot for identification. Four new plots were established at random locations within the grid system, bringing the total number of sampling plots to 44. In addition, existing plots were enlarged by moving the northwest corner rebar of each plot ten feet west, making all plot dimensions 1 foot x 20 feet.

Demographic plots were established at twenty-two sampling plots numbered 1, 2, 3, 4, 7, 8, 13, 15, 16, 19, 21, 23, 24, 25, 26, 33, 35, 38, 39, 40, 42, and 43. Due to intense ORV disturbance in 2003, several rebar posts were missing, including one or both corner rebar posts for plots 5, 28, 34, 38, and 39. Plots were reconstructed using the appropriate compass bearings and coordinates, and the rebar pieces were replaced.

2012-2019

In 2012, new monitoring and demography plots were established at this site because of vandalism to the plots in 2011. Sometime after population monitoring in 2011, most rebar posts that marked the long-term monitoring plot locations were removed making it impossible to relocate the original plots. The boundary of the population in French Flat South was mapped using a handheld GPSmap 60CSx, and a 157 m transect was laid at a bearing 180° due south to bisect the population. The head of the transect was marked with rebar capped with a yellow IAE cap and tagged with an aluminum disk embossed with #347. Additional rebar were placed at 50, 100 and 150m as well as at the end of the transect (157 m) and tagged with #348-351 respectively (Table 2, Figure 4, and Figure 6). Thirty-three density plots were run perpendicular to the baseline transect to the East or West at randomly selected intervals. A tape was run for 40 m and a 10 cm strip on the South side of the tape was monitored for *L. cookii*. Plants in the density plots were assigned to the life-history categories detailed below.

Density plots

We delineated plot boundaries with a meter tape and all plants within the plots were counted and assigned to a specific life-history category, as follows:

- S seedling
- V1/2 vegetative with 1 or 2 leaves
- V3 vegetative with 3 or more leaves
- R1 reproductive with 1 umbel
- R2 reproductive with 2 umbels
- R3 reproductive with 3 or more umbels

Life-history categories were originally developed for *L. bradshawii* monitoring in the Willamette Valley (Kaye et al. 2001). The similarities of the life-history characteristics of these species cause the categories to be applicable to *L. cookii* as well. Reproductive plants were segregated by umbel number because studies of *L. bradshawii* have shown that one-umbel plants rarely produce seed, while two-umbel plants produce seed on the second umbel, and three umbel plants may produce many seeds (Kaye 1992, Kaye and Kirkland 1994).

Table 2. Plot information for density and demography plots established in 2012 at French Flat South.

2012 New Plot #	Side of Tape	Location on Baseline (m)	End Rebar at (m)	End Rebar Tag	2012 Last Plant Found at (m)	Demog . Tag	Demog. Plot Location (NE Corner)	Demog. Plot Location (End)
362	E	13	23	363	9.3	-	-	-
364	W	27	30	365	10.4	-	-	-
366	W	30	33	367	6.9	-	-	-
707	W	36	21	708	15.9	329	10.5	11
709	W	38	21	710	15.8	330	6.5	7
711	W	42	21	712	12.7	331	4	4.5
749	E	45	40	750	23.6	353	15.5	15
713	E	52	37	714	25.8	332	17.5	17
741	E	57	35	742	27.1	352	20.5	20
743	E	59	40	744	30.1	354	21.5	21
753	E	61	36	754	31.8	357	23	22.5
745	W	65	39	746	23.9	361	13.5	14
747	E	70	40	748	32.1	360	8.5	8
751	E	72	40	752	22.8	355	5.5	5

LOMATIUM COOKII POPULATION MONITORING IN THE ILLINOIS VALLEY, JOSEPHINE COUNTY, OREGON

2012 New Plot #	Side of Tape	Location on Baseline (m)	End Rebar at (m)	End Rebar Tag	2012 Last Plant Found at (m)	Demog . Tag	Demog. Plot Location (NE Corner)	Demog. Plot Location (End)
725	W	79	40	726	11.8	338	3	3.5
715	E	81	40	716	30.4	333	11.5	11
717	W	94	40	718	28.7	334	13.5	14
719	E	95	40	720	30.9	335	19	18.5
721	E	97	40	722	30.6	336	15.5	15
723	W	99	32	724	31.0	337	16	16.5
727	W	107	28	728	24.6	339	13	13.5
701	E	109	40	702	27.2	326	24.5	24
703	E	111	40	704	28.3	546	23	22.5
705	W	116	32	706	23.0	328	19	19.5
729	W	119	33	730	22.8	340	24	24.5
755	E	125	40	756	27	356	16.5	16
731	E	126	40	732	19.1	341	5.5	5
733	W	128	35.5	734	29.2	342	20	20.5
735	E	129	40	736	23.6	343	9	8.5
737	W	136	40	738	31.5	344	15	15.5
757	W	142	33	758	32.8	358	5	5.5
759	W	144	34	760	32.6	359	9.5	10
739	W	154	40	740	32.8	345	18	18.5

LOMATIUM COOKII POPULATION MONITORING IN THE ILLINOIS VALLEY, JOSEPHINE COUNTY, OREGON




Figure 4. Layout of new density and demography plots established in 2012 at French Flat South and in 2013 at Middle. Density plots run perpendicular to the baseline to the east or west, and are monitored in 10 cm strips on the south side of the transect line. Demography plots are marked in the northeast and southwest corners (NE is tagged).




Figure 5. Location of French Flat Middle demographic and density plots established in 2013. Baseline transect is 100 m with a bearing of 034° (northeast). Last plants located on density plots are indicated by red crosses. Demography plots are indicated by blue squares and are not located along the baseline transect as shown here. See Table 1 for exact locations of all density and demography plots. Note that this figure does not reflect the transects on the east side of the baseline were extended in 2018.




Figure 6. Location of French Flat South demographic and density plots established in 2012. Baseline transect is 157 m with a bearing due south. Last plants located on density plots are indicated by red crosses. Demography plots are indicated by blue squares. See Table 2 for exact locations of all density and demography plots.

Demographic plots and analysis

Demographic plots were established at the Middle and South subpopulations in 1994. Sampling was suspended from 2000 through 2007 at the Middle subpopulation because sufficient information on population dynamics could be obtained from long-term study of one subpopulation to understand the basic life-history of the species and prepare a transition matrix model. In 2008-19, the Middle subpopulation was re-sampled and the updated information is presented below. Demographic monitoring at the South subpopulation has been consistent through the present year. In 2012 due to vandalism at French Flat South, new demographic plots were established (see below for details).

PLOT SAMPLING

At each subpopulation (Middle and South), 20 existing density plots were randomly selected as locations for 0.5 m x 0.5 m demographic plots (Table 1 and Table 2). Within each demographic plot, all *L. cookii* plants were mapped, given unique numbers (beginning with #1), assigned to the life history categories discussed earlier, and the presence or absence of grazing was recorded. To sample, a 0.5 meter x 0.5 meter frame was placed over the left rebar post, with the post positioned in the lower left corner of the frame. In 2012 at French Flat South the demographic plots were removed by vandals. We were not able to re-establish the plots, which had been monitored from 1993-2011, thus 30 new plots were monumented in 2012 (Table 2 and Figure 6). Location of the demographic plots were randomly selected along the randomly selected density transects. Two opposite corners of the 0.5 m x 0.5 m monitoring plot were marked with nails and the northeast corner is tagged (Figure 4).

CALCULATION OF SURVIVAL RATES AND ESTIMATES OF FERTILITY

All first-year vegetative plants with one or two leaves (and occasionally cotyledons) were considered seedlings; larger plants that appeared without previous observation were considered seedlings the previous year. We calculated the proportion of each stage that entered another stage (or remained the same) between consecutive years, and the proportion of individuals in each stage in each year. To estimate the number of seedlings produced by an individual in each reproductive stage (i.e., the fecundity of each stage), we used data on seed production from *L. cookii* sampled in 1996, the number of plants in each reproductive stage in the density plots, and the number of seedlings observed the following year in the density plots. These survival rates and estimates of fertility for each stage were arranged in 'transition matrices' for each pair of years from 1994-99 and 2008-19 for the Middle subpopulation and from 1994-2011 and 2012-19 for the South subpopulation. See Kaye and Pyke (2003), Kaye et al. (2001), Caswell and Kaye (2001), or Menges (1986) for complete discussions of transition matrix models for plant population dynamics and viability analyses.

POPULATION VIABILITY ANALYSIS – DETERMINISTIC AND STOCHASTIC MODELING

We evaluated population growth rate and viability with the transition matrices using data from the South subpopulation from 1994-2011 and 2012-19, and for the Middle subpopulation from 1994-99 and 2008-19. For this analysis, we calculated both deterministic and stochastic measures of population growth [λ , and stochastic λ] for each subpopulation. λ is the equilibrium population growth rate (and the dominant eigenvalue of the transition matrix), and can be used as a single measure of population viability to compare sites. Stochastic λ does not assume equilibrium population dynamics and incorporates observed environmental variability. If either type of growth rate is less than 1.0, the population will be projected to decrease in size, and eventually become extinct (a non-viable population). If λ is greater than 1.0, the population will grow (a viable population), given that

current conditions remain constant. For each site, we calculated λ for each year of observation. In addition, we used the average matrices from these available matrices to calculate an average population growth rate.

Assumptions of the model

Our use of the transition matrix model assumed that fertility and transition rates were independent of plant density. This is an acceptable assumption for many species with population densities below the density-dependent threshold (density-vague populations). However, density dependence eventually limits growth of populations with λ greater than one. Demographic stochasticity was also ignored by our models, but it usually generates little variation in population dynamics relative to environmental stochasticity, except at very low population sizes (Menges 1992). Our model assumed that population growth is a first-order Markov process, in which the probability that a plant will make a transition is independent of its stage in the previous year. In addition, we assumed that plants that first appeared as non-seedlings were not seedlings in the previous year, and that plants did not exhibit dormancy; years in which seedlings were not found or plants were skipped were considered missing data. We tested these assumptions with data collected through 2003, and they were found to have little effect on population growth rate compared to alternative assumptions (e.g., that new V2 and V3 plants were seedlings in the previous year, and that dormancy can occur).

Analysis

We conducted a population viability analysis incorporating environmental stochasticity. Population viability was evaluated in two different ways; stochastic population growth rate and extinction probability. This involved projecting future population dynamics by randomly selecting survival and fecundity measures from past years. We included environmental variability in our model through the matrix selection method. Matrix selection was accomplished by selecting a whole matrix at each time step, selected at random and with equal probability from the matrices available since demographic monitoring began in 1994. The matrices represent each year of the study, and the variation among them is environmental stochasticity. To calculate stochastic population growth rate, we used the program LAMS (Kaye 2001), converted to R script (R Core Team 2019), with 100,000 iterations.

We also calculated extinction probabilities for each subpopulation as a second measure of population viability. This simulation involved projecting future population dynamics by randomly selecting survival and fecundity measures from past years with the matrix selection method. More detailed descriptions of this method can be found elsewhere (i.e. Burgman et al. 1993, Kaye and Pyke 2002). These simulations ran for 20 years and consisted of 10,000 iterations. The starting population size for each simulation was 74,694 for the Middle subpopulation and 54,612 for the South subpopulation (the 1993 sizes), distributed among the six stages as follows: S=14%, V2=22%, V3=39%, R1=13%, R2=9%, and R3=3% for the South subpopulation and S=10%, V2=23%, V3=40%, R1=16%, R2=8%, and R3=3% for the Middle subpopulation. These proportions represent the average South and Middle subpopulation structures. The simulations stopped at the quasi-extinction threshold of 50% decline or 99% decline; this provided a conservative estimate of extinction dynamics. We used the program SHUFFLE (Kaye, unpublished program), written for MATLAB (Mathworks 1998) and converted to R (R Core Team 2019), to evaluate extinction probability with matrix selection methods.

In addition, we calculated elasticities of the mean matrix for both sites. Elasticities are the sensitivity of λ to small changes in the transition probabilities. Elasticities provide valuable information about

the extent to which population growth depends on survival, growth, and reproduction at different stages in the lifecycle (Caswell 1989). We used the statistical software R ver. 3.6.1 (R Core Team 2019) to calculate elasticities, as well as all other simulations and analyses.

Rough and Ready Creek

The population at Rough and Ready Creek was first documented in 1993 (Kagan 1994). In 1994, we established three transects for long-term monitoring of the two densest patches of *L. cookii* on BLM-managed land at this site (plants in the remaining patches were simply counted and the patches were mapped; Figure 8). To sample these transects, a 1 m x 1 m quadrat frame (constructed from light-weight $\frac{3}{4}$ inch PVC pipe) was placed on the ground at each meter mark along a measuring-tape stretched between rebar posts (**Error! Reference source not found.**). The individuals rooted within the frame were counted and assigned to the same stage categories used at French Flat.

Transects 1 and 2 are positioned within patch A (Figure 8). The origin for transect 1 was a rebar post set about 1 meter south of a flagged conifer and the endpoint was a rebar post set 15 meters away at a bearing of 130° east. The origin for transect 2 was a rebar post set 2 meters east of the origin-post for transect 1 and the endpoint was a rebar post set 15 meters away at a bearing of 130° east. Thus, transect 2 is adjacent and parallel to transect 1. [Note: In 1994, the origin for transect 2 was the same as used for the origin of transect 1; however, the plots along transect 2 began at the 3 meter mark to prevent overlap of plots 1, 2, and 3 along transect 1. The endpoint for transect 2 was a rebar post set 15 meters away at a bearing of 110° east from the origin post. This arrangement was altered in 1995 to reduce confusion.] Transect 3 (patch G) was located along the shoulder of the south side of the road across from transects 1 and 2. The origin rebar was located just above the shoulder of the road at a bearing of 150° east from the flagged conifer. This transect changed bearing at the 4.5 meter mark to accommodate the curve of the road. A second rebar post was set at 4.5 meters at a bearing of 120° east and the endpoint rebar was set at a bearing of 134° east and 7.8 meters from the second rebar post. Transect 3 parallels the shoulder of the road and passes through very dense patches of poison oak. In 2011 transect tapes were laid out along wheel ruts to define patches B, C, D, E and F.

In 2012 due to changes in plot set-up at French Flat South, all plants were monitored in a 10 cm strip along the south side of 40 m transects set-up perpendicular to the baseline transect. The starting location on the baseline transect is marked with rebar/PVC and tagged. The ends of the density transects are marked with tagged and capped rebar/PVC (Figure 7). The location along the baseline transect and the side of the baseline monitored (East or West) were determined using a random number generator. Thirty-three density transects were established in 2012.

Life-history categories were originally developed for *L. bradshawii* monitoring in the Willamette Valley (Kaye et al. 2001). The similarities of the life-history characteristics of these species cause the categories to be applicable to *L. cookii* as well. Reproductive plants were segregated by umbel number because studies of *L. bradshawii* have shown that one-umbel plants rarely produce seed, while two-umbel plants produce seed on the second umbel, and three umbel plants may produce many seeds (Kaye 1992, Kaye and Kirkland 1994).




Figure 7. Layout of new density and demography plots established in 2012 at French Flat South. Baseline transect is 157 m long. Density plots are run perpendicular to the baseline to the east or west. Density plots are monitored in 10 cm strips on the south side of the transect line. Demography plots are marked in the northeast and southwest corners (NE is tagged.) A table with locations of all density and demography plot locations can be found in Table 2.




Figure 8. Sketch map of the *L. cookii* patches located at Rough and Ready Botanical Wayside. Cross-hatching indicates patches of plants. The long-term monitoring transects are also indicated in the drawing. Note that the patches are distinguished by the letters A-H.

LOMATIUM COOKII POPULATION MONITORING IN THE ILLINOIS VALLEY, JOSEPHINE COUNTY, OREGON

Transects 1 and 2:

Transect line; plots oriented to either side of tape.

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
16	17	18	19	20	21	22	23	24	25	26	27	28	29	30

Transect 3:




Figure 9. Orientation of the sampling plots around transects 1, 2 and 3 at the Rough and Ready Botanical Wayside population.

Indian Hill

The population of *L. cookii* at Indian Hill was first documented in 1996, and monitoring was initiated in 1997. The population occurs on BLM managed land, but access is through private land. The population occupies a long stringer of open vegetation surrounded by dense forest, and extends to near the banks of the West Fork Illinois River.

In 1997, we established a grid system over the population. The perimeter of the population was carefully outlined to determine the extent of the plants and available habitat. The grid system had a bearing of 30° along its long axis for the first 100 meters then shifted westward to 5° for another 100 meters to conform to the shape of the habitat. Ten plots were sampled out of a total of 184 possible sample plot locations within the overall grid (Figure 11). All plots were 50 meters x 0.5 meters and were permanently marked at each end with rebar posts marked with flagging and copper tags with the plot number scratched onto the surface. To sample these plots, a 50 meter measuring tape was extended between rebar posts. The tape represented the west edge of the plot. Individual *L. cookii* plants that occurred within the plots (between the tape and 0.5 meter east) were counted and assigned to the same stage categories used at French Flat. Poison oak is patchy in this area, especially along the forest edges, so care should be exercised to minimize exposure. Also, *L. utriculatum* and *Perideridia* sp., which may resemble *L. cookii*, occur in this area and were encountered in the sampling plots (Figure 10).



Figure 10. *L. cookii* (left), and *L. utriculatum* (right). These species can co-occur in sampling plots.




Figure 11. Plot layout of the *Lomatium cookii* population at Indian Hill. Sample plots are 50 meters x 0.5 meters, marked at either end by a rebar post (with a copper tag and flagging). The plots are monitored on the east side of the tape.

RESULTS

French Flat

Population size and density

From the most recent population high of 137,560 in 2010, population size at the South subpopulation decreased 63% in 2019 to 50,811 (95% C.I.: 34,547 - 67,075), the lowest estimate of all sample years (Figure 12). Earlier increases in population size (from 1993 to 1998, Table 3) were likely due to several years of high seedling recruitment. The proportion of reproductive plants in the population declined from 1995 to 1997, reflecting the increased recruitment of seedlings that led to the total population increases observed in that time period (Figure 13 and Figure 14). This proportion peaked in 2008 at 37%, and steadily decreased to 11% in 2013, the lowest since 1997. Reproductive plant proportions had increased since 2013 to 33% in 2019 with a drop to 20% in 2017. In 2019, the density at the South subpopulation was 6.4 ± 1.0 (± 1 S.E.) plants m^{-2} , 38% of the long-term average. Density in this subpopulation has ranged from a low of 6.4 plants m^{-2} in 2019 to a high of 33 plants m^{-2} in 1997 and 1998 (Table 3).

While the South subpopulation decreased, the Middle subpopulation increased from 2018 to 2019 by 8.1% to 41,600 (95% C.I.: 27,091 - 56,109) (Figure 12). Population size grew from an initial 74,694 (95% C.I.: 53,954 - 95,433) in 1993 to a high of 195,057 (90% C.I.: 154,956 - 235,157) in 1998. The population decreased to its 2nd lowest size of 37,142 (24,181 - 50,102) in 2012 then increased to 81,067 (63,488 - 98,645) in 2013 and decreased to 39,933 (25,689 - 54,178) in 2014. The proportion of reproductive plants peaked at 41% in 2003 and 2008 but has steadily declined since 2008 (Figure 14). In 2019, the density at the middle subpopulation was 8.9 ± 1.6 (± 1 S.E.) plants m^{-2} , 44% of the long-term average. Density in this subpopulation ranged from a low of 5.8 plants m^{-2} in 2017 to a high of 41.7 plants m^{-2} in 1998 (Table 3).

Population structure

In 2019, the Middle and South subpopulations at French Flat were dominated by vegetative (V2-3) and small reproductive (R1) individuals (Figure 15). The large increase at Middle in 2013 (Table 3) was due to increased densities of vegetative plants from 5.6 and 5.1 m^{-2} in 2012 to 16.6 and 19.5 m^{-2} in 2013 for V1-2 and V3 plants respectively. On average, the vegetative classes were dominant, with V3 plants in the highest proportion. V3 plants have comprised the highest proportion for the South or Middle subpopulations since monitoring began, except for 1994, 1996-1998, and 2012.




Figure 12. The total number of *Lomatium cookii* at Middle and South French Flat, 1993-2019. Error bars represent a 95% confidence interval.




Figure 13. The total number of reproductive *Lomatium cookii* at Middle and South French Flat, 1993-2019. Error bars represent a 95% confidence interval.




Figure 14. The number of reproductive *Lomatium cookii* relative to the total subpopulation at Middle and South French Flat, 1993-2019.

Table 3. Summary of *Lomatium cookii* population data for the South and Middle French Flat subpopulations, 1993-2019. Population size and density estimates include all life-history stages from seedlings to large reproductive plants.

French Flat South		
Year	Population Size (95% CI)	Density (plant m⁻²) ± 1 SE
1993	54,612 (33,698 - 75,526)	6.8 ± 1.3
1994	164,837 (111,579 - 218,095)	20.6 ± 3.4
1995	169,350 (124,073 - 214,628)	21.2 ± 2.9
1996	247,452 (186,779 - 308,125)	31.0 ± 3.9
1997	264,524 (195,852 - 333,196)	33.1 ± 4.4
1998	267,075 (204,835 - 329,316)	33.4 ± 4.0
1999	239,014 (177,581 - 300,446)	29.9 ± 3.9
2000	202,416 (156,353 - 248,479)	25.3 ± 3.0
2001	174,943 (132,398 - 217,488)	21.9 ± 2.7
2002	174,551 (131,737 - 217,365)	21.8 ± 2.7
2003	154,829 (115,968 - 193,690)	19.4 ± 2.5
2004	113,227 (91,201 - 135,253)	14.2 ± 1.4
2005	123,432 (95,095 - 151,768)	15.4 ± 1.8
2006	98,215 (76,262 - 120,168)	12.3 ± 1.4
2007	80,260 (62,441 - 98,079)	10.0 ± 1.1
2008	109,008 (87,290 - 130,726)	13.6 ± 1.4
2009	129,907 (95,964 - 163,851)	16.3 ± 2.2
2010	137,560 (103,514 - 171,606)	17.2 ± 2.2
2011	127,454 (96,221 - 158,687)	16.0 ± 2.0
2012	122,650 (83,841 - 161,460)	15.4 ± 2.5
2013	114,296 (78,565 - 150,027)	14.3 ± 2.3
2014	74,379 (51,665 - 97,092)	9.3 ± 1.5
2015	59,565 (40,989 - 78,141)	7.5 ± 1.2
2016	81,640 (54,570 - 108,710)	10.2 ± 1.7
2017	53,282 (35,851 - 70,712)	6.7 ± 1.1
2018	63,847 (44,418 - 83,275)	8.0 ± 1.2
2019	50,811 (34,547 - 67,075)	6.4 ± 1.0

LOMATIUM COOKII POPULATION MONITORING IN THE ILLINOIS VALLEY, JOSEPHINE COUNTY, OREGON

French Flat Middle

Year	Population Size (95% CI)	Density (plant m ⁻²) ± 1 SE
1993	74,694 (53,954 - 95,433)	16.0 ± 2.3
1994	80,221 (56,402 - 104,040)	17.2 ± 2.6
1995	84,706 (61,621 - 107,790)	18.1 ± 2.5
1996	78,200 (60,376 - 96,023)	16.7 ± 2.0
1997	182,802 (144,335 - 221,269)	39.1 ± 4.2
1998	195,057 (154,956 - 235,157)	41.7 ± 4.4
1999	182,929 (145,214 - 220,643)	39.1 ± 4.1
2000	140,860 (112,090 - 169,631)	30.1 ± 3.2
2001	145,787 (115,652 - 175,922)	31.2 ± 3.3
2002	151,598 (121,965 - 181,232)	32.4 ± 3.3
2003	137,765 (106,531 - 168,999)	29.5 ± 3.4
2004	105,424 (80,107 - 130,741)	22.6 ± 2.8
2005	115,846 (88,950 - 142,743)	24.8 ± 3.0
2006	93,486 (68,163 - 118,809)	20.0 ± 2.8
2007	91,654 (66,336 - 116,971)	19.6 ± 2.8
2008	90,391 (70,338 - 110,443)	19.3 ± 2.2
2009	95,823 (72,378 - 119,268)	20.5 ± 2.6
2010	76,494 (55,832 - 97,156)	16.4 ± 2.3
2011	69,293 (52,478 - 86,109)	14.8 ± 1.8
2012	37,142 (24,181 - 50,102)	7.9 ± 1.4
2013	81,067 (63,488 - 98,645)	17.3 ± 1.9
2014	39,933 (25,689 - 54,178)	8.5 ± 1.6
2015	46,067 (32,665 - 59,468)	9.9 ± 1.5
2016	48,867 (37,243 - 60,490)	10.5 ± 1.3
2017	27,200 (17,917 - 36,483)	5.8 ± 1.0
2018	38,467 (23,510 - 53,423)	8.2 ± 1.6
2019	41,600 (27,091 - 56,109)	8.9 ± 1.6




Figure 15. *Lomatium cookii* population structure in 2019 (top) and averaged over the course of the study (bottom) for the Middle French Flat subpopulation (1994-2019), South French Flat subpopulation (1994-2019), Rough and Ready Creek (1995-2019), and Indian Hill (1997-2019). Bars indicate the mean percentage of plants in each stage. Error bars in the lower graph represent +1 SE. See text for explanation of life-history stages.

Demography and Population Viability Analysis

A total of 1,473 plants were mapped and categorized at the Middle subpopulation (1994-99: $n=745$ and 2008-19: $n=728$) and 1,241 at the South subpopulation (1994-2011: $n=826$ and 2012-19: $n=415$). Data from these 0.5 x 0.5 meter demographic plots were analyzed to construct transition matrices for Middle and South subpopulations (Table 4). In 2012, 286 new plants were mapped and categorized at the South subpopulation due to the loss of plot markers that year. Plants mapped in 2012 could not be identified to plants mapped prior to 2012, and thus there was not a 2011-12 transition matrix for the South subpopulation. New plots were established at the Middle subpopulation in 2013 to account for plots without plants for multiple years. Otherwise, plots prior to 2013 were identified, with exception of one, and annual transition matrices were constructed for 2012-19.

Deterministic growth rates at Middle and South have fluctuated since monitoring began, indicating that the French Flat *L. cookii* populations vary in population dynamics over space and time (**Error! Reference source not found.**). From 2008-17 at Middle, growth rates were below 0.9, with the lowest during 2016-17 of 0.587. Except for 2001 and 2006, growth rates at South have experienced a downward trend since 1995, when the population grew at its highest recorded rate of 1.522, and only two years, 2001 and 2006, had population growth rates above 1.0. The lowest population growth rate at South, 0.667, was observed in 2016-17. Growth rates at Middle and South in 2019 (0.877 and 0.795, respectively) were above and below the long-term average growth rates, respectively (Table 4).

In stochastic simulations incorporating environmental variability from 1994-2011 and 2012-19, South was projected to decline at an overall rate of 0.856. This reflects the overall poor annual growth rates at this site since monitoring began in 1994. At Middle, the stochastic growth rate was lower, 0.840, but this value represents observations from 1994-99 and 2008-19. The risk of sharp decline (50% loss) in 20 years was 99.9% at South and 99.7% at Middle. The risk of 99% decline in 20 years was 1.9% at the South subpopulation and 13.5% at the Middle subpopulation.

Thus, both deterministic and stochastic growth rates project that both South and Middle subpopulations will decline rapidly and substantially in the next two decades. The population growth rates and probabilities of decline reported here are best interpreted as comparisons of the performance of these two subpopulations and measures of their recent processes, rather than accurate predictions of future trends.

The elasticity analysis shows that small changes in the probability of V3 plants remaining in the V3 stage (0.183 at South and 0.231 at Middle) would have the single greatest effect on the population growth rate (Table 6). Overall, V3 plants had the highest cumulative elasticity (0.347 at South and 0.392 at Middle). The second highest cumulative elasticity was for the R2 stage at South (0.151) and the R1 stage at Middle (0.201). The seedling stage had the lowest cumulative elasticities (Table 6).

Annual mortality was lower in the Middle subpopulation than the South from 1994-98, except in 1995-96, and mortality was higher at Middle compared to South from 2009-2011 and 2012-2017 (**Error! Reference source not found.**). These mortality rates in the face of variable (sometimes positive) population growth over the same period indicate that the populations experience substantial dynamics, with individuals frequently changing size or dying with a variable influx of new plants. Complete information has not been tabulated on the life span of *L. cookii* individuals, but the information collected from the demographic plots can be used to determine or estimate how long individual plants live if the plots are sampled annually for several years. Of the 70 plants mapped at South as seedlings in 1994, 3 were still present in 2011 and 1 of them was reproductive. On average, close to half of seedlings

(50.7% at South and 57.0% at Middle) survive for two years. Only 2.7% on average observed seedlings became reproductive after one year in the South subpopulation during this entire study (1.4% in Middle). We found that, on average, 2.9% of seedlings became reproductive in 2 years, 6.0% in 3 years, and 5.9% in 4 years at the South subpopulation. The pattern of survival to reproductive status was similar at the Middle subpopulation (4.0% in 2 years, 6.9% in 3 years, and 11.8% in 4 years).

Table 4. Average transition matrices used in the stochastic analyses for *L. cookii* at French Flat. The top row of each matrix represents the number of seedlings produced per reproductive plant (fecundity). Lower rows represent the probability of an individual in a given stage changing to another stage the following year. Stage-specific mortality is provided in the bottom row. The elements in the matrices are averaged from the transition data collected in 1994-2011 and 2012-19 for the South subpopulation and 1994-99 and 2008-19 for the Middle subpopulation.

South, average matrix (1994-2011 & 2012-19), $\lambda = 0.871$						
	S	V2	V3	R1	R2	R3
S	0	0	0	0	0.456	2.839
V2	0.418	0.295	0.134	0.066	0.062	0.037
V3	0.258	0.379	0.460	0.236	0.162	0.078
R1	0.023	0.035	0.123	0.252	0.234	0.197
R2	0.018	0.008	0.056	0.175	0.224	0.156
R3	0	0.004	0.008	0.020	0.095	0.322
Mortality	0.284	0.278	0.220	0.251	0.223	0.210
Middle, average matrix (1994-99 & 2008-19), $\lambda = 0.863$						
	S	V2	V3	R1	R2	R3
S	0	0	0	0	0.630	1.828
V2	0.469	0.354	0.135	0.056	0.067	0
V3	0.177	0.288	0.509	0.312	0.119	0.218
R1	0.004	0.041	0.128	0.320	0.321	0.278
R2	0.005	0.019	0.033	0.140	0.198	0.163
R3	0	0.006	0.009	0.026	0.079	0.205
Mortality	0.345	0.292	0.187	0.147	0.216	0.137

Table 5. Population growth and mortality rates at the French Flat Middle and South sub-populations from 1994-2019.
[Note: Demographic data was not collected at the Middle subpopulation from 2000 through 2007, and plants mapped prior to 2012 were not relocated at the South subpopulation.]

year	Lambda		Mortality (%)	
	Middle	South	Middle	South
1994-95	1.103	1.142	18%	39%
1995-96	1.554	1.522	17%	15%
1996-97	1.016	0.811	11%	32%
1997-98	1.135	0.931	10%	25%
1998-99	0.900	0.692	12%	30%
1999-00	N/A	0.789	N/A	18%
2000-01	N/A	0.771	N/A	26%
2001-02	N/A	1.114	N/A	24%
2002-03	N/A	0.687	N/A	35%
2003-04	N/A	0.837	N/A	23%
2004-05	N/A	0.830	N/A	33%
2005-06	N/A	0.933	N/A	20%
2006-07	N/A	1.011	N/A	32%
2007-08	N/A	0.906	N/A	18%
2008-09	0.873	0.855	21%	28%
2009-10	0.844	0.993	25%	22%
2010-11	0.816	0.846	27%	22%
2011-12	0.617	N/A	36%	N/A
2012-13	0.681	0.803	34%	24%
2013-14	0.719	0.838	29%	17%
2014-15	0.617	0.724	43%	32%
2015-16	0.821	0.816	20%	17%
2016-17	0.587	0.667	37%	36%
2017-18	0.986	0.990	9%	17%
2018-19	0.877	0.795	16%	23%

Table 6. Elasticity matrices of the south and middle *Lomatium cookii* subpopulations derived from the average transition matrices. Elasticities represent the relative sensitivity of the population growth rate (λ) to small changes in the transition probabilities. The sum of all elasticities is 1.0, and the sum of each column is the total proportional sensitivity of λ to changes in transition probabilities in that stage. The top two individual elasticities are shown in bold.

South subpopulation

	S	V2	V3	R1	R2	R3
S	0	0	0	0	0.029	0.058
V2	0.044	0.051	0.042	0.007	0.004	0.001
V3	0.034	0.082	0.183	0.032	0.014	0.002
R1	0.004	0.009	0.059	0.042	0.024	0.006
R2	0.005	0.004	0.046	0.049	0.039	0.009
R3	0	0.004	0.017	0.014	0.041	0.045
sum	0.087	0.150	0.347	0.144	0.151	0.121

Middle subpopulation

	S	V2	V3	R1	R2	R3
S	0	0	0	0	0.03	0.029
V2	0.038	0.068	0.047	0.008	0.004	0
V3	0.019	0.071	0.231	0.057	0.009	0.005
R1	0.001	0.013	0.074	0.075	0.030	0.009
R2	0.001	0.009	0.027	0.047	0.027	0.007
R3	0	0.005	0.013	0.015	0.018	0.016
sum	0.058	0.165	0.392	0.201	0.118	0.065

Rough and Ready Creek

Due to its relatively small size we censused the entire *L. cookii* population at Rough and Ready Creek. In 2019, we counted 2,444 *L. cookii* plants in eight patches at Rough and Ready Creek (**Error! Reference source not found.**, Figure 16, and Figure 17), a 50% increase from 2018. While there was year-to-year stochasticity as well as variation among patches, overall, the population has increased since monitoring began.

The proportion of reproductive plants in the population in 2019 (39%) was lower than the long-term average of 45% (Figure 16). Small and large vegetative plants (V2-3) and small reproductive plants (R1) were the most abundant stages at Rough and Ready Creek in 2019 (Figure 15). And on average since 1994, the proportion of large reproductive plants (R2-3) was higher at Rough and Ready Creek than at any other population (Figure 15).

LOMATIUM COOKII POPULATION MONITORING IN THE ILLINOIS VALLEY, JOSEPHINE COUNTY, OREGON

Table 7. Number of *L. cookii* individuals in each patch at Rough and Ready Creek. The total number of plants in all patches combined represents a census of the population. Totals include plants in all stages from seedlings to large reproductive plants. In 2015, patches C, D, E, and F were lumped into patch B, and in 2018, patch E was lumped into F.

Year	Number of plants per patch								Total
	A	B	C	D	E	F	G	H	
1994	144	6	3	15	13	10	80	7	278
1995	177	79	22	85	127	53	135	14	692
1996	259	82	44	82	30	42	119	14	670
1997	167	56	104	315	91	131	119	12	990
1998	141	82	93	180	128	48	106	7	785
1999	223	237	201	308	393	197	81	14	1,654
2000	228	167	105	171	102	296	61	11	1,141
2001	292	113	102	189	245	119	64	24	1,148
2002	88	153	104	135	274	217	16	18	1,005
2003	298	93	217	64	34	382	73	25	1,186
2004	251	114	123	46	271	174	74	25	1,078
2005	248	146	124	56	148	369	72	25	1,188
2006	98	271	235	69	424	540	50	28	1,715
2007	127	141	67	132	217	197	23	20	924
2008	397	250	477	157	433	325	79	9	2,127
2009	567	363	430	424	347	526	102	17	2,776
2010	346	567	758	300	133	382	60	11	2,557
2011	480	394	389	404	406	671	71	15	2,830
2012	292	349	486	146	336	489	65	16	2,179
2013	360	432	393	188	141	392	58	9	1,973
2014	300	450	377	238	263	175	15	9	1,827
2015	497	1,522	-	-	-	-	126	13	2,158
2016	405	300	383	142	198	261	114	15	1,818
2017	233	69	636	61	69	440	117	17	1,642
2018	131	107	724	76	-	452	127	13	1,630
2019	360	342	1,164	46	12	384	130	6	2,444




Figure 16. Population trends and the total number of plants (reproductive and vegetative) at Rough and Ready Creek, 1994-2019. There were only 26 reproductive plants recorded in 1994.




Figure 17. Population fluctuations in *L. cookii* patches A-H at Rough and Ready Creek, 1994-2019. Patches B-F were combined in 2015 and patches E-F were combined in 2018.

Indian Hill

The total population size of *L. cookii* at Indian Hill has generally increased since monitoring was initiated in 1997. There were approximately 166 more plants in 2019 than in 2018, for a total of 8,225 (Figure 18 and **Error! Reference source not found.**). Population size in 2016 (5,943) was the second lowest on record behind the size in 2000 (5,704). On average, reproductive plants have composed 32% of the population (minimum: 16% in 1997; maximum: 46% in 2003). In 2019, reproductive plants were 44% of the total population. The 1997-1998, 1999-2000, and 2015-2016 declines in total population size are the result of attrition of vegetative plants, while the increase in density from 2002-2005 and 2007-2011 can be attributed to a rise in the proportion of reproductive plants. The mean density of reproductive plants in the population increased steadily from 0.4 plants m^{-2} in 1997 to 0.81 plants m^{-2} in 2004, 2005, and 2006. The density of reproductive plants in 2010 was 0.87 plants m^{-2} , the highest since monitoring began. In 2019, there was approximately 0.78 reproductive plants m^{-2} , an increase of 29% from the previous year.

Most plants at Indian Hill were vegetative (Figure 18). In 2019, the V2 and V3 stages composed 26% and 29% of the population, respectively. The R1 stage also made a substantial contribution (34%) to the population in 2019, while all reproductive stages totaled 44% of the population. Seedling numbers at the site have varied substantially, from 0% in 1999, 2000, and 2014 to 16% in 2005. The proportion of seedlings decreased from 3.9% in 2018 to 1.6% in 2019.




Figure 18. Number of vegetative and reproductive plants at Indian Hill, 1997-2019.

Table 8. Summary of population density (mean $\pm$ 1 S.E.) and size (mean $\pm$ 95% confidence limit) for the Indian Hill population, 1997-2019.

year	density (m ⁻²)		population size	
	reproductive	total	reproductive	total
1997	0.40 $\pm$ 0.10	2.56 $\pm$ 0.65	1,840 $\pm$ 913	11,776 $\pm$ 5,738
1998	0.46 $\pm$ 0.21	1.41 $\pm$ 0.51	2,134 $\pm$ 1,842	6,477 $\pm$ 4,466
1999	0.39 $\pm$ 0.15	1.54 $\pm$ 0.49	1,803 $\pm$ 1,335	7,084 $\pm$ 4,279
2000	0.46 $\pm$ 0.19	1.24 $\pm$ 0.45	2,098 $\pm$ 1,674	5,704 $\pm$ 3,904
2001	0.58 $\pm$ 0.29	1.54 $\pm$ 0.64	2,686 $\pm$ 2,562	7,084 $\pm$ 5,599
2002	0.60 $\pm$ 0.27	1.41 $\pm$ 0.54	2,778 $\pm$ 2,367	6,477 $\pm$ 4,698
2003	0.73 $\pm$ 0.30	1.58 $\pm$ 0.75	3,349 $\pm$ 2,668	7,250 $\pm$ 6,556
2004	0.82 $\pm$ 0.39	1.91 $\pm$ 0.83	3,772 $\pm$ 3,387	8,795 $\pm$ 7,259
2005	0.81 $\pm$ 0.32	2.80 $\pm$ 1.35	3,735 $\pm$ 2,826	12,898 $\pm$ 11,874
2006	0.80 $\pm$ 0.30	2.14 $\pm$ 0.90	3,680 $\pm$ 2,660	9,862 $\pm$ 7,849
2007	0.48 $\pm$ 0.22	1.89 $\pm$ 1.13	2,226 $\pm$ 1,969	8,685 $\pm$ 9,942
2008	0.60 $\pm$ 0.27	2.04 $\pm$ 1.04	2,870 $\pm$ 2,374	9,623 $\pm$ 9,018
2009	0.76 $\pm$ 0.31	2.17 $\pm$ 0.99	3,478 $\pm$ 2,721	9,973 $\pm$ 8,640
2010	0.87 $\pm$ 0.48	2.38 $\pm$ 1.28	3,993 $\pm$ 4,186	10,948 $\pm$ 11,263
2011	0.70 $\pm$ 0.36	2.48 $\pm$ 1.23	3,220 $\pm$ 3,161	11,426 $\pm$ 10,784
2012	0.36 $\pm$ 0.15	2.02 $\pm$ 0.78	1,638 $\pm$ 1,281	9,310 $\pm$ 6,822
2013	0.54 $\pm$ 0.26	2.29 $\pm$ 1.04	2,502 $\pm$ 2,306	10,543 $\pm$ 9,077
2014	0.56 $\pm$ 0.25	1.95 $\pm$ 0.77	2,558 $\pm$ 2,156	8,979 $\pm$ 6,711
2015	0.48 $\pm$ 0.23	1.86 $\pm$ 0.68	2,190 $\pm$ 1,999	8,538 $\pm$ 5,949
2016	0.36 $\pm$ 0.10	1.29 $\pm$ 0.38	1,974 $\pm$ 912	5,943 $\pm$ 3,374
2017	0.56 $\pm$ 0.35	1.57 $\pm$ 0.96	2,558 $\pm$ 1,545	7,213 $\pm$ 4,298
2018	0.60 $\pm$ 0.25	1.75 $\pm$ 0.69	2,778 $\pm$ 2,215	8,059 $\pm$ 6,046
2019	0.78 $\pm$ 0.30	1.79 $\pm$ 0.60	3,588 $\pm$ 2,647	8,225 $\pm$ 5,253

Herbivory on *Lomatium cookii*

Herbivory by voles is relatively frequent on *L. cookii*, but the frequency of damage to the plants has varied substantially among sites and years. Typically, damage to the plants ranges from consumption of leaf tips and inflorescences, to removal of nearly all aboveground plant material. Consumption of the flower clusters results in little damage to the individual plant, but eliminates seed production and, therefore, recruitment of seedlings in the following year. The long-term impact of herbivory on *L. cookii* plants is unknown, but appears to play an important role in population growth and seedling dynamics. Environmental parameters (e.g., rainfall, average maximum daily temperature, and average minimum daily temperature) may influence vole populations similarly across the region and in turn affect the number of *L. cookii* that are grazed.

Herbivory on *L. cookii* has fluctuated between years, and the patterns are not synchronous among the sites (Figure 19). Average herbivory at French Flat Middle and South increased since initial monitoring. In 2019, French Flat South and Rough and Ready populations had decreased herbivory from 2018, while French Flat Middle and Indian Hill populations had increased herbivory in the same year. In our observations since 1993, the highest recorded average herbivory was 57% (95% C.I.: 46% - 68%) detected at Indian Hill in 2019, while the lowest was 1% (95% C.I.: 0% - 2%) at French Flat Middle in 1994.

LOMATIUM COOKII POPULATION MONITORING IN THE ILLINOIS VALLEY, JOSEPHINE COUNTY, OREGON




Figure 19. Frequency of herbivory on *L. cookii* populations at four populations, 1994-2019. Error bars represent 95% confidence intervals. All plants were measured at Rough and Ready population.

DISCUSSION

Current population status

In 2019, the number of *L. cookii* increased at all sites except for French Flat South.

French Flat

Current population estimates at this site are approximately 41,600 plants at the Middle subpopulation and 50,811 at the South subpopulation. The population viability analysis suggests that the South and Middle subpopulations may be at risk of long term decline if current conditions persist. ORV use remains frequent and damaging (Figure 20). IAE is analyzing the effects of climate on *L. cookii* through Climate Driven Population Models and the results will be included in a separate report.

The Middle subpopulation hosted an estimated 195,057 individuals in 1998 when the population was at its highest, up from 74,694 when monitoring began in 1993 and over seven times the current estimate. Although there has been some variability, the population has generally been declining since 2000. The South subpopulation also peaked in 1998, with a total of 267,075 individuals. Since then, it has generally declined as well.

Results from population viability analyses based on information from 1994-99 and 2008-19 for Middle and 1994-2011 and 2012-19 for South indicated that these subpopulations differed in their long-term growth rates by about 2%, although both are likely to decline. Stochastic simulations based on observed environmental variability indicated that the risk of extinction or catastrophic decline was high at the Middle and South subpopulation if past environmental and biological conditions persist.

Rough and Ready Creek

We counted 2,444 plants at Rough and Ready Creek in 2019, which was a 50% increase from 2018. The Rough and Ready Creek population occurs in grassy patches and roadways on gentle to moderate slopes among shrubs and trees (*Pinus jeffreyi*). This patchy, less extensive habitat pattern is probably responsible for the large difference in population size between this site and French Flat. While road use can reduce the number of plants in this population, we observed recruitment along the roadbeds in 1995-2006, likely due to the reduced competition in these areas. A large trash pile found on the western edge of Patch A in 2003 and fresh ORV tracks in multiple years are indicative of active road use and off-road habitat impact. Boulders placed to block vehicle traffic on some roads and grassland patches has resulted in effective protection of this habitat, and grasses are now colonizing the protected road beds.

Indian Hill

The Indian Hill population, which is intermediate in size between the Rough and Ready and French Flat populations, declined in total numbers from 1997 (when monitoring began) to 1998, then remained relatively stable from 1998 to 2004. Overall, the population has increased slightly and remained stable from 2005-19, although 2016 had the second lowest population size on record. Estimates of total plant density and reproductive plant density followed similar changes. Many of the V3 and R1 plants are likely a result of the prolific seedling recruitment in 2005 and 2011. The proportion of seedlings in the population has been increasing since a low point in 2014. The population size has been relatively stable in recent years, and the site does not appear to receive ORV use. The error around our estimates of plot density and population size at Indian Hill have increased since monitoring began. This is due to high variability among transects and increased shrub density from *Ceanothus cuneatus*.



Figure 20. ORV damage in the *Lomatium cookii* population at French Flat.

Management Recommendations

French Flat

We recommend that population monitoring at the French Flat subpopulations be conducted once per year for all plots. If insufficient time is available for these activities, sampling of the density plots should be conducted at least once every three years for data collection and plot maintenance. Sampling of the subpopulation boundary transects can occur at five-year intervals (or when changes in population boundaries are suspected). Additional data from the demographic plots will allow improved population viability analyses, including enhancement of environmental stochasticity estimates and greater certainty in evaluations of the importance of environmental variables such as precipitation and vole herbivory.

Maintaining the motorized vehicle closure should be a priority for management of this site. Off Road Vehicles still use this area, despite official closure of the roads, making this important population

extremely vulnerable to damage (Figure 20). Also, horse-back riders that use French Flat should be warned of the presence of rebar posts interspersed throughout the habitat.

Prescribed burning was implemented at French Flat South and Middle in 2015 to allow for comparisons of population processes before and after burning. In 2017, density of *L. cookii* did not differ between burned and unburned plots or among vegetation/reproductive class (Giles et al. 2017).

Rough and Ready Creek

The road leading through the Rough and Ready population is un-gated but some sensitive portions of habitat at this site have been blocked off from vehicle traffic with boulders. These obstacles appear to be having the desired effect of blocking vehicle access, which was a significant concern at this site in recent years. Vehicle access should be monitored at this site to confirm that the boulders remain in place and that habitat protection is succeeding.

Monitoring at Rough and Ready Creek should be repeated each year to detect changes in population size and density. However, if insufficient resources are available for this activity, sampling should be conducted at least once every three years.

Indian Hill

The habitat at Indian Hill is restricted by forest that surrounds a long strip of meadow habitat. These meadows are being invaded by shrubs that may be having detrimental impacts on the *L. cookii* population through competition for light, water, and nutrients. We highly recommend that these shrubs be removed to improve habitat conditions. Dense shrub cover is impeding abilities to accurately assess population size and we have seen decreases in population size; removal of shrubs adjacent to the population is recommended. Monitoring at Indian Hill should be repeated annually to detect changes in population size and density.

LITERATURE CITED

- Caswell, H. and T.N. Kaye. 2001. Stochastic demography and conservation of an endangered perennial plant (*Lomatium bradshawii*) in a dynamic fire regime. *Advances in Ecological Research* 32:1-51.
- Ferson, S. 1991. RAMAS/stage: generalized stage-based modeling for population dynamics. Applied Biomathematics. Setauket, New York.
- Giles, D.E.L., E.C. Gray, and M.A. Bahm. 2017. Assessing management techniques for *Lomatium cookii* (Cook's desert-parsley). Institute for Applied Ecology, Corvallis, Oregon and USDI Bureau of Land Management, Medford District.
- Kagan, J. 1980. The biology of *Lomatium bradshawii* (Apiaceae), a rare plant of Oregon. M.S. Thesis, University of Oregon, Eugene, OR.
- Kagan, J. 1986. A new species of *Lomatium* (Apiaceae), from southwestern Oregon. *Madroño* 33: 71-75.
- Kagan, J. 1994. Habitat management plan for *Lomatium cookii* (Cook's desert-parsley) in the Illinois Valley, Josephine County, OR. Oregon Natural Heritage Program. Portland, OR.
- Kaye, T.N. 1992. Bradshaw's desert-parsley: population monitoring and pollination biology. *Kalmiopsis* 2:1-4.
- Kaye, T.N. 2001. Population viability analysis of Bradshaw's lomatium (*Lomatium bradshawii*) and a comparison of matrix modeling techniques. PhD Dissertation. Department of Botany and Plant Pathology, Oregon State University, Corvallis, OR.
- Kaye, T.N. and M. Blakeley-Smith. 2002. Vegetation survey of French Flat ACEC, Medford District, BLM. Institute for Applied Ecology, Corvallis, Oregon and USDI Bureau of Land Management, Medford District.
- Kaye, T.N., K. Pendergrass, K. Findley and J.B. Kauffman. 2001. The effect of fire on the population viability of an endangered prairie plant. *Ecological Applications* 11:1366-1380.
- Kaye, T.N. and M. Kirkland. 1994. Population biology of *Lomatium bradshawii*. II. Sexual expression, breeding system, and insect interactions. Unpub. Cooperative Challenge Cost Share report, Oregon Department of Agriculture, Plant Conservation Biology Program and Eugene District, BLM.
- Kaye, T.N. and M. Kirkland. 1995. *Lomatium cookii* population monitoring at French Flat and Rough and Ready Botanical Wayside, Illinois Valley, Josephine County, Oregon. Unpub. Challenge Cost Share report, Oregon Department of Agriculture, Plant Conservation Biology Program and Medford District BLM.
- Kaye, T.N. and D.A. Pyke. 2003. The effect of stochastic technique on estimates of population viability from transition matrix models. *Ecology* 84:1464-1476.
- Knight, L. 1992. Baseline monitoring of *Lomatium cookii*. Bureau of Land Management, Medford District. Medford, Oregon.
- Mathworks. 1998. Matlab, version 5. The Mathworks, Inc., Natick, Massachusetts.
- Menges, E.S. 1986. Predicting the future of rare plant populations: demographic monitoring and modeling. *Natural Areas Journal* 6:6-17.

LOMATIUM COOKII POPULATION MONITORING IN THE ILLINOIS VALLEY, JOSEPHINE COUNTY, OREGON

- R Core Team. 2019. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org/>.
- Tong, B. 1993. *Lomatium cookii* monitoring at French Flat, Freeman claim. Unpublished Medford District BLM staff report.

APPENDIX A. GEAR LIST.

Field Gear

Last year's report

Last year's datasheets

Density plot datasheets

French Flat South and Middle demographic datasheets

Rite-in-the-rain paper

Jepson guide

Clipboards/pencils

Maps

3 100+ foot tapes, 2 smaller (50ft) tapes

candy canes

rulers

large binder clips

Bundle of pin flags

Quadrat frames: 1x1m for Rough and Ready, .5x.5m for French Flat

Extra rebar, hammer, wire, tags and flagging to replace lost/bent

Compass

First Aid Kit/Tecnu

Water jug *2

Pruning tool

Health and Safety box

Note for planning:

- Indian Hill – 10 transects took about 4-5 hours. Park at turn-off for dirt road, unless conditions have been improved since 2010. May need to walk in from gate on Waldo Rd.
- Need vehicle with clearance for Indian Hill.